Originally published as:

Tibi, R., Wiens, D. A., Shiobara, H., Sugioka, H., Yuan, X. (2007): Double seismic discontinuities at the base of the mantle transition zone near the Mariana slab. - *Geophysical Research Letters*, 34, L16316

DOI: [10.1029/2007GL030527](https://doi.org/10.1029/2007GL030527).

1 Double Seismic Discontinuities at the Base of the Mantle
2 Transition Zone near the Mariana Slab

3 Rigobert Tibi¹, Douglas A. Wiens¹, Hajime Shiobara², Hiroko
4 Sugioka³, and Xiaohui Yuan⁴

5 ¹ *Department of Earth and Planetary Sciences, Washington University,*
6 *St. Louis, Missouri, USA.*

7 phone: 314 935 7965

8 fax: 314 935 7361

9 email: tibi@seismo.wustl.edu

10 ² *Earthquake Research Institute, University of Tokyo, Tokyo, Japan.*

11 ³ *Institute for Frontier Research on Earth Evolution, Japan Agency for Marine-Earth*
12 *Science Technology, Yokosuka, Japan*

13 ⁴ *GeoForschungsZentrum Potsdam, Potsdam, Germany*

14 submitted to *Geophysical Research Letters*, April 28, 2007

15 revised July 27, 2007

16 Abstract

17 We use P -to- S converted phases from teleseismic data recorded at island and ocean
18 bottom stations in Mariana to investigate the upper mantle structure in the region.
19 We find evidence for double seismic discontinuities at the base of the transition zone
20 near the Mariana slab. A shallower discontinuity is imaged at depths of ~ 650 – 715 km
21 and a deeper interface lies at ~ 740 – 770 km depth. The large lateral extent at near
22 constant depths for both features is consistent with horizontal interfaces rather than
23 small-scale scatterers. The amplitude ratios of the seismic signals suggest that the
24 shear velocity contrast across the two interfaces is comparable. These characteristics
25 support the notion that the discontinuities are the results of phase transformations in
26 olivine (ringwoodite to post-spinel) and non-olivine component (ilmenite to perovskite),
27 respectively, for the pyrolite model of mantle composition.

1. Introduction

The 660-km seismic discontinuity marking the bottom of the mantle transition zone is believed to result from an isochemical phase transition involving olivine, namely the dissociation of ringwoodite into perovskite plus magnesiowüstite [Ringwood, 1969]. However, mineral physics calculations and high-pressure experiments suggest that, depending upon the bulk mantle composition, transformations of non-olivine components such as the garnet system may lead to a complex structure of the 660-km discontinuity [Vacher *et al.*, 1998; Hirose, 2002]. This is the case particularly along cold temperature profiles, similar to those expected in subduction zone environments. Under cold conditions, the transformation of garnet to ilmenite, and the transition from ilmenite to perovskite should cause two additional discontinuities near the base of the transition zone. These interfaces and the discontinuity induced by ringwoodite dissociation are expected to occur at distinct depths, causing multiple interfaces [Vacher *et al.*, 1998].

Multiple seismic discontinuities near the base of the transition zone have been detected in thermally complex regions such as beneath southern California [Simmons and Gurrola, 2000] and northeast China, at the tip of the subducting Japan slab [Niu and Kawakatsu, 1996; Ai *et al.*, 2003]. The 660-km discontinuity is routinely observed in global studies of long-period *SS* precursors [Shearer, 1993; Gossler and Kind, 1996]. Its non-detection on a global scale using precursors to long-period *PP* waves [Estabrook and Kind, 1996; Shearer and Flanagan, 1999] has recently been attributed to a complicated and highly variable structure, that shows single and double reflections, consistent with multiple phase transitions [Deuss *et al.*, 2006].

We applied the *P*-wave receiver-function methodology [e.g., Vinnik, 1977] on teleseismic data recorded at island and ocean bottom stations in Mariana to investigate the upper mantle structure in the region and found evidence for double seismic discontinuities at the base of the transition zone around the Mariana slab.

2. Data and Analysis

Data used in this study are seismograms from 315 earthquakes with epicentral distances from 30 to 95°, recorded at a temporary network in the Mariana region (Fig. 1). The network, which consists of 21 broadband island stations and 27 semi-broadband ocean bottom seismographs (OBS) operated from June 2003 to May 2004 [Tibi *et al.*, 2006]. Also included are data recorded between 1991 and 2005 at the permanent station GUMO, located on the island of Guam.

The *P*-wave receiver-function technique applied in this study is well established and described in greater detail in Vinnik [1977] and Yuan *et al.* [1997]. Only key steps of the methodology are mentioned here. For land stations, the three-component seismograms are deconvolved to ground displacement and filtered at 2–30 sec. In order to isolate *P*-to-*SV* converted phases, the seismograms are rotated to the LQT ray coordinate system. The earthquake source-time function is removed by deconvolving the *P* energy on the L component from the rotated traces. The Q component contains *P*-to-*SV* conversions. For the seafloor data, oceanic noise was successfully suppressed with an 8–30 sec bandpass filter. The orientations of the OBS's were determined using polarization angles obtained from airgun shots and large, well-located, distant events [Pozgay, pers. comm.]. The subsequent rotation and deconvolution procedures were performed the same way as for land data. We computed a total of 972 high-quality Q component receiver functions (RF's).

In order to enhance low-amplitude *Ps* converted phases from the upper mantle velocity discontinuities, we divided the study area into 1°×2°-boxes and stacked moveout-corrected RF's having piercing points at 660 km depth within the same box. Only stacks resulting from a minimum of 10 individual RF's were used to search for discontinuity conversions. The dimensions of the boxes were chosen so that there are sufficient RF's for each box, while insuring that the discontinuity phases within each box illuminate nearly the same structure. In a subduction zone, the most likely cause of major variations in the structure of the upper mantle discontinuities is the differ-

ence in temperature between the subducting plate and the neighbouring mantle. The north-south orientation of the Mariana slab implies that these variations would occur primarily in the east-west direction. Hence, the dimension of the boxes in that direction were chosen to be only half the dimension in the north-south direction. For each box, we estimated the 95% confidence levels of the RF stack by performing bootstrap resampling [Efron and Tibshirani, 1991]. Only phases in the stack for which the lower confidence limit was clearly above zero were interpreted. This insures that random noise in the data are not misinterpreted as conversion from the discontinuities. Because of the difference in their frequency content (Fig. 2a and d), land and OBS recordings were stacked separately. However, for the few boxes with stacks both from land and OBS data, the inferred discontinuity depths were consistent among the two datasets within the estimated range of uncertainty.

3. Results and Discussion

Moveout-corrected RF's from land stations for one of the $1^\circ \times 2^\circ$ -boxes is shown in Fig. 2a. The trace to the right is the stack of the RF's. In order to distinguish deep P_s conversions from multiples originating from shallow structures, RF's are stacked in slowness bins of 0.2 sec/deg to improve the signal to noise ratio, and subsequently sorted in order of increasing mean slowness. The procedure aligns P_s phases parallel to the slowness axis, while multiples become more inclined (Fig. 2b). The phases labeled '410', '660' and '750' can be correlated across nearly the entire slowness range at about 42, 68 and 74 sec, respectively, suggesting that these phases are direct P_s conversions. In contrast, the delay time of the phase labeled 'mul.' decreases with increasing slowness from about 53 sec to 48 sec, as expected for multiple reverberations (Fig. 2b). This interpretation is corroborated by the fact that this phase is aligned parallel to the slowness axis and enhanced in the stack, when the traces are moveout-corrected for multiple reverberations $PpPs$ (Fig. 2c). The phase, which is observed throughout the study area, is interpreted as $PpPs$ from an interface at about 150 km depth in the mantle wedge. The phases at about 42 and 68 sec in Fig. 2a are

110 conversions from the 410-km and 660-km discontinuity at depths of about 390 and 675
 111 km, respectively. We interpret the converted energy at ~ 74 sec as originating from an
 112 interface at about 740 km depth (hereinafter referred to as the 750-km discontinuity).
 113 The low-frequency stack of the OBS RF's displayed in Fig. 2d shows basically similar
 114 features as the land data. In this case, however, the deep interface is observed at ~ 760
 115 km depth, while conversion from the 660-km discontinuity does not rise significantly
 116 above the background noise level.

117 Following *Kind et al.* [2002] we migrated the RF's obtained from land stations using
 118 a modified IASP91 velocity model [*Kennett and Engdahl*, 1991]. The modification
 119 included a crustal P velocity model for the Mariana arc obtained from an active source
 120 study [*Takahashi et al.*, 2007]. The crustal S velocities were derived from that model
 121 using a V_P/V_S of 1.73. Fig. 3 shows an east-west section of the migrated traces
 122 along the AB profile indicated in Fig. 1. It includes RF's having piercing points at
 123 660 km depth located within the latitude range of 15–17°N. The 660-km discontinuity
 124 is observed intermittently from longitude about 144.5 to 147.5°E, and appears to be
 125 depressed to ~ 705 km depth at 145.5°E. The location of the depression coincides
 126 with the likely deep projection of the slab, suggesting that it is induced by the cold
 127 slab temperatures, as expected for the negative Clapeyron slope of the ringwoodite
 128 transition to post-spinel [*Ito and Takahashi*, 1989]. The lateral extent of only ~ 100
 129 km for the depression suggests that in this region the Mariana slab penetrates the
 130 660-km discontinuity with little or no advective thickening. Conversions from the
 131 interface at ~ 750 km depth are observed between longitude about 145 and 148°E. The
 132 intermittence of the observations is principally due to gaps in the interface sampling.

133 Results of the box-by-box stacking are summarized in Fig. 4. We estimate the
 134 uncertainty in discontinuity depths to be ± 10 km, coming mainly from picking errors
 135 and uncertainties of the velocity model associated with the slab anomaly. The 660-
 136 km discontinuity is observed in the depth range between about 650 and 715 km, with
 137 the greatest depths occurring in the latitude range of 15–17°N within or near the
 138 Mariana slab (Fig. 4a). The inferred maximum depth for the 660-km discontinuity is

well within the range of 710–730 km obtained from a high-resolution study of P_{660p} and S_{660p} phases recorded at regional stations [Tibi *et al.*, 2006]. The topography of the 750-km interface shown in Fig. 4b indicates a variation between about 740 and 775 km in depth. For most of the boxes in which we detected the 750-km interface, we also observed the 660-km discontinuity (Fig. 4). This indicates the existence of two distinct discontinuities, demonstrating that the deeper interface is not the 660-km discontinuity that has migrated to greater depth. Whereas the section in Fig. 3 shows double discontinuities between 145 and 148°E, in Fig. 4 around 16°N, double discontinuities are only seen between 146 and 147°E. Variations in the structure of the 750-km interface across the relatively large stacking boxes in Fig. 4 are the most likely cause for this discrepancy.

Earlier studies have detected small-scale scatterers near the top of the lower mantle (at depths between 710 and 900 km) around the Mariana slab [Krüger *et al.*, 2001; Kaneshima, 2003]. The scatterers located within a high velocity anomaly [Bijwaard *et al.*, 1998] have been interpreted as a piece of lithosphere or fragments of former oceanic crust subducted below the 660-km discontinuity [Krüger *et al.*, 2001; Kaneshima, 2003]. The relatively large lateral extent of the 750-km feature and its near constant depth throughout the study area (Fig. 3 and 4b) are clearly consistent with a horizontal interface rather than a series of small-scale scatterers, which would be randomly located. For each stack showing conversion both from the 660-km and 750-km discontinuity, we calculated the amplitude ratio between the two phases (P_{750s}/P_{660s}). The amplitudes of P -to- S conversion depend primarily on the magnitude of shear velocity contrast across the converting interface. Considering that P_{750s} and P_{660s} travel similar paths, and that differential attenuation due to the different conversion depths is negligible, the inferred average value of 1.01 ± 0.22 for P_{750s}/P_{660s} indicates that the shear velocity contrast across the two discontinuities is comparable.

The most likely causes of deep, flat interfaces are isobaric phase transformations or compositional changes. Mineral physics calculations by Vacher *et al.* [1998] suggest that for a pyrolite model of mantle composition, at temperature conditions similar to

those expected in subduction zones, garnet transforms to ilmenite within the depth range of 608–664 km. Under these conditions, the breakdown of ringwoodite takes place at depths of 690–693 km, while ilmenite transforms to perovskite at 709–731 km depth. According to these authors, the garnet to ilmenite transition occurs in a very broad pressure range, inducing a discontinuity that is rather weak. This may be the reason why we found no evidence for such interface. In contrast, both the breakdown of ringwoodite and the transformation of ilmenite to perovskite are expected to cause relatively strong discontinuities of similar sharpnesses. Our observations appear to be consistent with these predictions. The occurrence of the 750-km discontinuity seems to be restricted to regions surrounding the slab (Fig. 3 and 4b), assumed to be pyrolite mantle, further supporting our interpretation.

An alternative model for the observed 750-km interface is that it may be the result of chemical layering associated with metasomatism in the uppermost lower mantle. In that model, the transport of elements by fluids released from descending slab may cause chemical layering or heterogeneity in terms of $\text{Mg}_2\text{O}/\text{SiO}_2$ ratios [Ohtani *et al.*, 2001]. As the Mariana slab is near-vertical, slab-expelled fluids could rise through the neighbouring mantle west as well as east of the slab, explaining the observation of the 750-km discontinuity both on the mantle wedge side and trenchward behind the slab (Fig. 3 and 4b).

Acknowledgments. Equipment for the land-based stations was provided by the PASSCAL program of IRIS. Lamont-Doherty provided the US OBS's. We are grateful to Megan Flanagan and an anonymous referee for their helpful review. This research was financially supported by NSF under grants OCE0001938 and EAR0310272.

References

- 192
193 Ai, Y., T. Zheng, W. Zu, Y. He, and D. Dong (2003), A complex 660 km discontinuity
194 beneath northeast China, *Earth Planet. Sci. Lett.*, *212*, 63–71.
- 195
196 Bijwaard, H., W. Spakman, and E. R. Engdahl (1998), Closing the gap between re-
197 gional and global travel time tomography, *J. Geophys. Res.*, *103*, 30,055–30,078.
- 198
199 Deuss, A., S. A. T. Redfern, K. Chambers, and J. H. Woodhouse (2006), The nature
200 of the 660-kilometer discontinuity in Earth’s mantle from global seismic observations of
201 *PP* precursors, *Science*, *311*, 198–201.
- 202
203 Efron, B., and R. Tibshirani (1991), Statistical data analysis in the computer age,
204 *Science*, *253*, 390–395.
- 205
206 Engdahl, E. R., R. van der Hilst, and R. Buland (1998), Global teleseismic earth-
207 quake relocation with improved travel times and procedures for depth determination,
208 *Bull. Seismol. Soc. Am.*, *88*, 722–743.
- 209
210 Estabrook, C. H., and R. Kind (1996), The nature of the 660-kilometer upper-mantle
211 seismic discontinuity from precursors to the *PP* phase, *Science*, *274*, 1179–1182.
- 212
213 Gossler, J., and R. Kind (1996), Seismic evidence for very deep roots of continents,
214 *Earth Planet. Sci. Lett.*, *138*, 1–13.
- 215
216 Hirose, K. (2002), Phase transitions in pyrolitic mantle around 670-km depth: Im-
217 plications for upwelling of plumes from the lower mantle, *J. Geophys. Res.*, *107*,
218 10.1029/2001JB000597.
- 219

- 220 Ito, E., and E. Takahashi (1989), Postspinel transformations in the system Mg_2SiO_4 -
 221 Fe_2SiO_4 and some geophysical implications, *J. Geophys. Res.*, *94*, 10,637–10,646.
- 222
- 223 Kaneshima, S. (2003), Small-scale heterogeneity at the top of the lower mantle around
 224 the Mariana slab, *Earth Planet. Sci. Lett.*, *209*, 85–101.
- 225
- 226 Kennett, B. L. N., and E. R. Engdahl (1991), Traveltimes for global earthquake loca-
 227 tion and phase identification, *Geophys. J. Int.*, *105*, 429–465.
- 228
- 229 Kind, R. et al. (2002), Seismic images of crust and upper mantle beneath Tibet:
 230 Evidence for Eurasian plate subduction, *Science*, *298*, 1219–1221.
- 231
- 232 Krüger, F., M. Baumann, F. Scherbaum, and M. Weber (2001), Mid mantle scat-
 233 terers near the Mariana slab detected with a double array method, *Geophys. Res.*
 234 *Lett.*, *28*, 667–670.
- 235
- 236 Niu, F., and H. Kawakatsu (1996), Complex structure of mantle discontinuities at
 237 the tip of the subducting slab beneath northeast China: A preliminary investigation
 238 of broadband receiver functions, *J. Phys. Earth*, *44*, 701–711.
- 239
- 240 Ohtani, E., M. Toma, K. Litasov, T. Kubo, and A. Suzuki (2001), Stability of dense
 241 hydrous magnesium silicate phases and water storage capacity in the transition zone
 242 and lower mantle, *Phys. Earth Planet. Inter.*, *124*, 105–117.
- 243
- 244 Ringwood, A. E. (1969), Phase transformations in mantle, *Earth Planet. Sci. Lett.*, *5*,
 245 401–412.
- 246
- 247 Shearer, P. M. (1993), Global mapping of upper mantle reflectors from long-period
 248 *SS* precursors, *Geophys. J. Int.*, *115*, 878–904.

249

250 Shearer, P. M., and M. P. Flanagan (1999), Seismic velocity and density jumps across
 251 the 410- and 660-kilometer discontinuities, *Science*, *285*, 1545–1548.

252

253 Simmons, N. A., and H. Gurrola (2000), Multiple seismic discontinuities near the base
 254 of the transition zone in the Earth’s mantle, *Nature*, *405*, 559–562.

255

256 Takahashi, N. et al. (2007), Crustal structure and evolution of the Mariana intra-
 257 oceanic island arc, *Geology*, *35*, 203–206.

258

259 Tibi, R., D. A. Wiens, H. Shiobara, H. Sugioka, and P. J. Shore (2006), Depth of
 260 the 660-km discontinuity near the Mariana slab from an array of ocean bottom seis-
 261 mographs, *Geophys. Res. Lett.*, *33*, L02313, doi:10.1029/2005GL024523.

262

263 Vacher, P., A. Mocquet, and C. Sotin (1998), Computations of seismic profiles from
 264 mineral physics: The importance of the non-olivine components for explaining the 660
 265 km depth discontinuity, *Phys. Earth Planet. Inter.*, *106*, 275–298.

266

267 Vinnik, L. P. (1977), Detection of waves converted from P to SV in the mantle, *Phys.*
 268 *Earth Planet. Inter.*, *15*, 294–303.

269

270 Yuan, X., J. Ni, R. Kind, J. Mechie, and E. Sandvol (1997), Lithospheric and up-
 271 per mantle structure of southern Tibet from a seismological passive source experiment,
 272 *J. Geophys. Res.*, *102*, 27,491–27,500.

Figure Captions

Fig. 1. Bathymetric map of the Mariana region. Squares and triangles are locations of island stations and OBS sites, respectively. Hexagon indicates the station GUMO of the Global Seismic Network. Stars represent deep (depth >300 km) seismicity. The earthquakes include events re-located by *Engdahl et al.* [1998], and best located earthquakes that occurred during the operational period of the Mariana network. Ray-theoretical P s piercing points at 660 km depth for the calculated receiver functions are represented with crosses and dots for land and OBS data, respectively. The white line indicates the location of the profile displayed in Fig. 3. The inset at the upper-right corner shows the locations of the 315 events used in this study (circles), and the Mariana seismic network (triangle).

Fig. 2. (a) Moveout-corrected receiver functions. The traces were corrected for the directly converted phase P s and a reference slowness of 6.4 sec/deg using a modified IASP91 velocity model [*Kennett and Engdahl*, 1991]. Receiver functions are from island stations, and include data having P s piercing points at 660 km depth located within the box at 11–13°N and 146–147°E. Delay time is relative to the direct P wave. The thin horizontal lines indicate the predicted arrival times for P -to- S conversions at 410 and 660 km depth. The trace to the right is the stack of the receiver functions for this box. The gray areas in the stack indicate robust conversions for which the lower 95% confidence limit is above zero. Conversions from the 410, 660-km, and 750-km discontinuities are marked with arrows. Multiples from an interface at about 150 km depth are labeled by ‘mul.’ (see text). (b) Slowness-stacked receiver functions. The waveforms shown in (a) have been averaged over slowness bins of 0.2 sec/deg. The slowness range of 7–9 sec/deg corresponds to the range of epicentral distances of about 31–58° for the events that sample the box. Horizontal dotted lines at about 42, 68 and 74 sec delay time indicate conversions from the 410, 660 and 750-km discontinuities, respectively. White dots mark multiples from the 150-km interface. (c) The same as in

301 (b) except that the traces were moveout-corrected for multiple reverberations $PpPs$.
 302 (d) The same as in (a) for the OBS data and the box at 15–17°N and 142–143°E.

303

304 **Fig. 3.** East-west profile of migrated receiver function data. The location of the
 305 profile is shown in Fig. 1. Data are from island stations, and include receiver functions
 306 with Ps piercing points at 660 km depth located within the latitude range of 15–17°N.
 307 Red (blue) indicates velocity increase (decrease) downward. The black thick arrow
 308 indicates the position of the slab, as defined by seismicity (dots). The thin horizontal
 309 line indicates the depth of 660 km. Conversions from the 660-km discontinuity and
 310 750-km interface are marked with black and green line segments, respectively. Note
 311 the small-scale depression between 145 and 146°E for the 660-km discontinuity.

312

313 **Fig. 4.** Maps of the discontinuity depth. Crosses and dots indicate Ps piercing
 314 points at 660 km depth for land and OBS data, respectively. Circles represent deep
 315 (depth >300 km) seismicity. The discontinuity depth for each $1^\circ \times 2^\circ$ -box was estimated
 316 from the stack obtained after summation of the RF's having Ps piercing points at 660
 317 km depth located within the box. Boxes with weak conversions from the discontinuities
 318 or null observations are indicated in white. (a) Depths of the 660-km discontinuity.
 319 (b) The same as in (a) for the 750-km discontinuity.

Figure 1: Tibi *et al.*

Figure 2: Tibi *et al.*

Figure 3: Tibi *et al.*

Figure 4: Tibi *et al.*