



Originally published as:

Łuszczak, K., Persano, C., Braun, J., Stuart, F. M. (2017): How local crustal thermal properties influence the amount of denudation derived from low-temperature thermochronometry. - *Geology*, 45, 9, pp. 779–782.

DOI: <http://doi.org/10.1130/G39036.1>

# How local crustal thermal properties influence the amount of denudation derived from low-temperature thermochronometry

Katarzyna Łuszczak<sup>1</sup>, Cristina Persano<sup>1</sup>, Jean Braun<sup>2</sup>, and Finlay M. Stuart<sup>3</sup>

<sup>1</sup>School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ, UK

<sup>2</sup>Helmholtz Centre Potsdam, German Research Center for Geosciences (GFZ), Telegrafenberg, 14473 Potsdam, Germany

<sup>3</sup>Isotope Geoscience Unit, Scottish Universities Environmental Research Centre (SUERC), East Kilbride G75 0QF, UK

## ABSTRACT

Translating amounts and rates of rock cooling derived from low-temperature thermochronometry into denudation requires assumptions about the local geothermal gradient. The temperature gradient in the crust depends on many factors, including basal heat flow, crustal heat production, and thermal conductivity. Consequently, geothermal gradients may be variable on time scales over which rock cooling is tracked by thermochronometry. Using one-dimensional numerical modeling of heat transfer in rocks of varying thermal characteristics, we show that the geothermal gradient of the eroded layer is the most important factor for accurate estimation of denudation amounts. Using a three-dimensional numerical model (Pecube), we demonstrate the impact of crustal heat production and thermal conductivity on estimates of total denudation derived from apatite fission track data from central west Britain. We show that the regional variation in cooling ages measured in Caledonian granites can be explained by geothermal gradient variation due to the presence of a heat-producing granite batholith and removal of insulating sedimentary rocks, and does not require variable denudation. Neglecting the blanketing effect leads to twofold overestimation of the amount of denudation. The occurrence of heat-producing basement that was once covered by a sedimentary blanket is common, in particular in the core of mountain belts. Accurate determination of the amount and rate of denudation from thermochronometric studies in these situations must take into account the composition of the eroded rocks.

## INTRODUCTION

Low-temperature thermochronometry (LTT) is used to quantify amounts of denudation by converting paleotemperatures into paleodepths. In the absence of a clear indication of elevated basal heat flow, present-day geothermal gradients (20–30 °C/km; Turcotte and Schubert, 2002) are typically assumed. This assumption ignores the spatial and temporal variation in heat production and thermal conductivity of rocks, as well as short-lived heat-flow perturbations, such as from faulting, fluid circulation, magmatism, and rapid rock uplift (Turcotte and Schubert, 2002). Although efforts have been made to predict the effect of these factors on the upper crustal geothermal gradient (e.g., Brown et al., 1994), only recently have studies considered the role of the thermal conductivity of the rocks that are being denuded (e.g., Barbarand et al., 2013; Braun et al. 2016), while variation in crustal heat production is largely neglected.

Surface heat flow is the sum of basal heating of the lithosphere and radiogenic heat production in the crust. Typically, each contributes about equally to surface heat flow (Turcotte and Schubert, 2002). However, where the upper crust comprises high-heat-producing (>3 μW/m<sup>3</sup>) rocks, crustal heat production may contribute in excess of 80% (Neumann et al., 2000). The

thermal conductivity in the uppermost crust varies from 0.5 to 7.0 W/m/K depending on temperature, pressure, porosity, quartz content, and water saturation (Eppelbaum et al., 2014). Low-conductivity rocks such as mudstone, coal, and basalt have an insulating effect on the underlying rocks (often referred to as the “blanketing effect”) that can enhance the maturity of organic material in sedimentary basins (Pollack and Cercone, 1994) and the geothermal energy potential of heat-producing basement (Majorowicz and Minea, 2012).

Acid-intermediate basement rocks are commonly sampled for LTT due to the abundance of apatite and zircon. Exhumation events commonly involve erosion of overlying sedimentary or volcanic successions. In such cases, the combined effect of the low-conductivity rocks overlying heat-producing igneous bodies on denudation estimates derived from LTT data has not yet been investigated. Here we use one-dimensional (1-D) and 3-D (Pecube) numerical models to determine the extent to which the thermal conductivity of eroded strata and heat production in the basement affect the geothermal gradient in the uppermost crust. The modeling results are tested in a case study from central west Britain, where we show that denudation can be overestimated by a factor of two if variations in the thermal structure of the crust are ignored.

## QUANTIFYING GEOTHERMAL GRADIENTS

Heat transfer in eroding crust follows a second-order equation:

$$\frac{\partial T}{\partial t} + v \frac{\partial T}{\partial z} = \kappa \frac{\partial^2 T}{\partial x^2} + \kappa \frac{\partial^2 T}{\partial y^2} + \kappa \frac{\partial^2 T}{\partial z^2} + A, \quad (1)$$


where  $T$  is temperature,  $t$  is time,  $v$  is velocity of rock uplift,  $x$ ,  $y$ , and  $z$  are coordinates of the rock particle,  $\kappa$  is thermal diffusivity, and  $A$  is rate of heat production ( $\kappa = k/\rho c$  and  $A = H/\rho c$ , where  $k$  is thermal conductivity,  $H$  is heat production,  $\rho$  is density, and  $c$  is specific heat capacity). In Figure 1 we show solutions to the 1-D (vertical) form of this equation for scenarios designed to determine how the geothermal gradient changes in response to heat production in a pluton and to the thermal conductivity of an overlying sediment pile.

Figure 1A shows how geothermal gradients change in response to pluton thickness and heat production (constant thermal conductivity). For a 4–12-km-thick intrusive body, the geothermal gradient increases only by 1.2–4.4 °C/km for each 1 μW/m<sup>3</sup> increase in  $H$ .

For a “normal” crust, where heat production is low and constant (1 μW/m<sup>3</sup>), the geothermal gradient increases rapidly when the thermal conductivity of the blanketing layer is <2 W/m/K (Fig. 1B); high-porosity–low-conductivity rocks, such as chalk (≤1.5 W/m/K; Eppelbaum et al., 2014), can produce geothermal gradients of >40 °C/km.

Figure 1C shows the combined effect on the geothermal gradient of varying radiogenic heat production and thermal conductivity. The two factors can either produce extremely low or high geothermal gradients, or almost cancel each other, leaving the gradient practically unchanged. The point to note here is that geothermal gradients in excess of 50 °C/km can readily be generated in a low-conductivity sedimentary layer overlying a high-heat-producing granite.

Figure 1D shows the thermal structure of the crust for varying heat production and with or without a sedimentary cover. For 2 km of erosion, calculating denudation using the geothermal gradient in the preserved rocks is accurate only when the sedimentary cover is absent. In all other cases, denudation is significantly




**Figure 1.** A–C: Plots showing effect of heat production ( $H$ ) and thermal conductivity ( $k$ ) on geothermal gradient in uppermost crust made of: heat-producing body of variable  $H$  and thickness,  $k = 2.5 \text{ W/m/K}$  (constant), intruded in “normal” crust (A); 1-km-thick sedimentary layer with variable  $k$  and  $H = 0 \mu\text{W/m}^3$  (constant), overlying “normal” crust (B); or 1-km-thick sedimentary layer with variable  $k$  (values in figure) and  $H = 0 \mu\text{W/m}^3$  overlying 12-km-thick body ( $H$  varies,  $k = 2.5 \text{ W/m/K}$ ) (C). D: Change of temperature with depth (black lines) for different crustal compositions (GS—granite with sedimentary cover; CS—“normal” crust with sedimentary cover; UC—uniform “normal” crust; G0—granite only;  $z$ —elevation relative to paleosurface). Granite is 12 km thick with  $H = 5 \mu\text{W/m}^3$  and  $k = 2.5 \text{ W/m/K}$ . Blanket layer is 1 km thick with  $k = 1.5 \text{ W/m/K}$  and  $H = 0 \mu\text{W/m}^3$ . Shaded area represents rocks preserved after 2 km of denudation. Gray lines indicate predicted denudation if blanket layer is not accounted for. In all models, “normal” crust has  $k = 2.5 \text{ W/m/K}$  and  $H = 1 \mu\text{W/m}^3$ .

overestimated, demonstrating that an accurate estimate of the geothermal gradient within the eroded rock section is crucial for accurate determination of the amount of denudation using LTT.

### CASE STUDY—CENTRAL WEST BRITAIN

Rapid exhumation of onshore Britain in the early Paleogene is consistent with rock uplift in response to the arrival of the Iceland mantle plume; however, the amount and extent of denudation is controversial (Green et al., 2012; Cogné et al., 2016). Apatite fission track (AFT) data show that latest Cretaceous temperatures in the English Lake District were  $>110 \text{ }^\circ\text{C}$  and were  $50\text{--}90 \text{ }^\circ\text{C}$  in the surrounding regions (Fig. 2B), requiring  $>3 \text{ km}$  of denudation centered on the  $\sim 60\text{-km}$ -wide Lake District block (assuming a geothermal gradient of  $30 \text{ }^\circ\text{C/km}$ ; Green, 1986). This amount of exhumation is at odds with stratigraphic reconstructions (Holliday, 1993) and estimates based on the thickness of magmatic underplating (Tiley et al., 2004). An elevated geothermal gradient ( $\sim 60 \text{ }^\circ\text{C/km}$ ; Green, 2002) was introduced to reconcile these conflicting interpretations and was explained as an effect of enhanced basal heat flow from the underplating melts (Green et al., 2012). However, deep-seated magmatism has a negligible effect on the thermal structure of shallow crust (see Fig. DR2 in the GSA Data Repository<sup>1</sup>; Brown et al., 1994). Holliday (1993) proposed that the

<sup>1</sup>GSA Data Repository item 2017256, the apatite fission track data set and modeling parameters, is available online at <http://www.geosociety.org/datarepository/2017/> or on request from [editing@geosociety.org](mailto:editing@geosociety.org).



**Figure 2.** Maps of Great Britain showing: subsurface extent of granite batholiths and sampling locations (A); latest Cretaceous temperatures (circles show data from Green et al. [1997], Thomson et al. [1999], and Green [2002]; stars show data from this study) (B); and surface heat flow, after Busby et al. (2011) (C).

high latest Cretaceous temperatures were a consequence of high heat flow and a now-eroded low-conductivity sedimentary cover. This interpretation has remained untested.


We collected new AFT data from the Lake District, south Scotland, and north Wales to explore the effect that a low-conductivity sedimentary cover and variable heat production may have on cooling histories (Fig. 2A). New AFT ages are youngest in the Lake District (48–75 Ma) and significantly older (>150 Ma) in south Scotland and north Wales. Thermal histories were derived using a Bayesian approach of data inversion in the QTQt software (Gallagher, 2012). The models indicate that latest Cretaceous temperatures were ~120 °C in the Lake District and ~70 °C in south Scotland and north Wales (Fig. 2B), similar to those presented by Green et al. (1997). The full data set and paleotemperatures estimates are available in the Data Repository.

The latest Cretaceous temperature pattern roughly correlates with the present-day surface heat flow (Fig. 2C; Busby et al., 2011). Heat flow exceeds 80 mW/m<sup>2</sup> around the large granite batholiths in north England, where radiogenic heat production is 1.9–5.2 μW/m<sup>3</sup> (Downing and Gray, 1986). These values are two to five times the average heat production in Phanerozoic crust (0.9–1.1 μW/m<sup>3</sup>; Jaupart and Marschal, 2005). The AFT-derived thermal histories

indicate an exhumation pulse that began in the latest Cretaceous, when the granites were overlain by Mesozoic mudstones and chalk (Holiday, 1993). The low thermal conductivity of these rock types (1.2–2.0 W/m/K; Downing and Gray, 1986) implies that they may have provided thermal insulation to enhance the temperatures in the granites at the time of denudation.

### Effect of Thermally Heterogeneous Crust


In order to test the effect of variable heat production and sedimentary cover on the AFT-derived thermal histories, three sets of forward models (grand total of 56) for three crust thermal structures were determined using the Pecube numerical model (Braun et al., 2012): UC, a spatially uniform crust with a geothermal gradient of 20–30 °C/km; G0, a spatially uniform crust that includes four heat-producing bodies (one underneath the Lake District and three in south Scotland); and GS, a crust structure similar to G0, but covered by a blanket of low-conductivity sedimentary rocks. Models were run with variable thermal parameters and for different values of total rock uplift. The quality of the models is evaluated based on the misfit ( $\mu$ ) between the observed and predicted AFT ages. All modeling parameters, misfit values obtained for particular models, and a detailed description of the modeling procedure are given in Section DR-2c of the



**Figure 3.** Misfit of Pecube forward models versus amount of total uplift for three analyzed scenarios: UC—uniform crust with geothermal gradient of 20–30 °C/km; G0—crust with heat-producing granite; GS—crust with granite and sedimentary cover. Yellow shaded area shows range of uplift values acceptable by regional geological constraints.

Data Repository. Misfit values are plotted versus total rock uplift in Figure 3. For a range of total uplift consistent with regional geological constraints, low misfit values (<5) are found only by models that consider the combined effect of high-heat-producing granites and a sedimentary blanket.

Figure 4 shows, for each modeled scenario, the predicted distributions of AFT ages for total rock uplift of 2.25 km, which, for a given topography, implies exhumation of 1.25–2.25 km. Measured AFT ages are significantly younger



**Figure 4.** Pecube-derived forward models of distribution of apatite fission track (AFT) ages for total uplift of 2.25 km and different crustal compositions (UC—uniform crust; G0—crust with heat-producing granite; GS—crust with granite and sedimentary cover) for study area in Great Britain (see Fig. 2 for location). Lower graphs show predicted versus observed ages for sampled localities. Gray line is 1:1 reference line. Panel on left (Topo) shows digital elevation model, location of granite batholiths (dashed lines), and range of thermal parameters used for batholiths and sedimentary blanket:  $r$ —radius of batholith;  $A$ —heat production rate (LD—Lake District granite; Scot—Scottish granites);  $K$ —thermal diffusivity.

and more variable than those that the uniform crust model (UC) generates. The AFT age distribution is better reproduced if a heat-producing body is present (G0), but the modeled ages are still older than measured values. The best fit to the data ( $\mu < 4$ ) is obtained when, prior to exhumation, a low-conductivity sedimentary succession covers a heat-producing intrusion (GS). In this scenario, the AFT age pattern is the result of variable thermal properties of the rocks, not variable denudation.

The model has some limitations, such as a simplified spatial distribution of thermal parameters and a lack of pre-Cenozoic uplift and subsidence events. Although taking these aspects into account would further improve our estimates of denudation, they have little effect on the regional pattern of the AFT ages on which the model results are based (see Section DR-2d in the Data Repository).

The 3-D numerical models support the results of the 1-D model by demonstrating that crustal thermal heterogeneities have to be taken into consideration when deriving the amount of uplift and denudation from LTT data. In this case, the presence of a low-thermal-conductivity layer reduces the amount of total Cenozoic denudation required in the Lake District by 50%–70%, and the AFT-derived cooling is translated into 1.25–2.25 km of denudation. These amounts overlap with the stratigraphic estimates of eroded Mesozoic cover (1.2–1.75 km; Holliday, 1993), without the need to invoke elevated basal heat flow.

## IMPLICATIONS

This study demonstrates that not accounting for the increased geothermal gradient in a low-thermal-conductivity sedimentary layer underlain by high-heat-producing rocks may lead to significant overestimation of the amount of denudation. This issue can be particularly important in LTT studies, as granitic rocks are favored because of their content of idiomorphic apatite and zircons and they are commonly exhumed by erosion of overlying sedimentary successions. For instance, in the European Alps, the Variscan granite batholiths have been exposed by post-orogenic erosion, subsequently buried beneath Paleozoic–Paleogene sediments, and finally exhumed during Alpine orogenesis, resulting in complex LTT age patterns (e.g., Glotzbach et al., 2011) that are likely influenced by the thermal properties of the eroded crust. Similar situations can be found in other orogenic belts (Carrapa et al., 2014; Hoke et al., 2015), at rifted and passive margins (Persano et al., 2006; Wildman et al., 2015), and in intra-plate settings (Majorowicz and Minea, 2012; Barbarand et al., 2013). This study shows that quantifying the amount and regional extent of denudation should be preceded by a careful examination of the thermal properties of both sampled and eroded rocks.

## ACKNOWLEDGMENTS

The funding for this study was provided by the University of Glasgow, UK, (the College of Science and Engineering Ph.D. Scholarship and the Sir Alwyn Williams Scholarship) and the Natural Environment Research Council (NERC) standard grant (GU 52562-NE/H008276/1). Kerry Gallagher is thanked for providing the code for the 1-D modeling. Andy Gleadow, Zoltán Erdős, and one anonymous reviewer are thanked for their helpful and constructive comments.

## REFERENCES CITED

- Barbarand, J., Quesnel, F., and Pagel, M., 2013, Lower Paleogene denudation of Upper Cretaceous cover of the Morvan Massif and southeastern Paris Basin (France) revealed by AFT thermochronology and constrained by stratigraphy and paleosurfaces: *Tectonophysics*, v. 608, p. 1310–1327, doi:10.1016/j.tecto.2013.06.011.
- Braun, J., van der Beek, P., Valla, P., Robert, X., Herman, F., Glotzbach, C., Pedersen, V., Perry, C., Simon-Labric, T., and Prigent, C., 2012, Quantifying rates of landscape evolution and tectonic processes by thermochronology and numerical modeling of crustal heat transport using PECUBE: *Tectonophysics*, v. 524–525, p. 1–28, doi:10.1016/j.tecto.2011.12.035.
- Braun, J., Stippich, C., and Glasmacher, U.A., 2016, The effect of variability in rock thermal conductivity on exhumation rate estimates from thermochronological data: *Tectonophysics*, v. 690, p. 288–297, doi:10.1016/j.tecto.2016.09.027.
- Brown, R., Gallagher, K., and Duane, M., 1994, A quantitative assessment of the effects of magmatism on the thermal history of the Karoo sedimentary sequence: *Journal of African Earth Sciences*, v. 18, p. 227–243, doi:10.1016/0899-5362(94)90007-8.
- Busby, J., Kingdon, A., and Williams, J., 2011, The measured shallow temperature field in Britain: *Quarterly Journal of Engineering Geology and Hydrogeology*, v. 44, p. 373–387, doi:10.1144/1470-9236/10-049.
- Carrapa, B., Orme, D.A., DeCelles, P.G., Kapp, P., Cosca, M.A., and Waldrup, R., 2014, Miocene burial and exhumation of the India-Asia collision zone in southern Tibet: Response to slab dynamics and erosion: *Geology*, v. 42, p. 443–446, doi:10.1130/G35350.1.
- Cogné, N., Doepke, D., Chew, D., Stuart, F.M., and Mark, C., 2016, Measuring plume-related exhumation of the British Isles in Early Cenozoic times: *Earth and Planetary Science Letters*, v. 456, p. 1–15, doi:10.1016/j.epsl.2016.09.053.
- Downing, R.A., and Gray, D.A., 1986, *Geothermal Energy: The Potential in the United Kingdom*: London, British Geological Survey, 187 p.
- Eppelbaum, L., Kutasov, I., and Pilchin, A., 2014, *Applied Geothermics*: Berlin, Heidelberg, Springer, 751 p., doi:10.1007/978-3-642-34023-9.
- Gallagher, K., 2012, Transdimensional inverse thermal history modeling for quantitative thermochronology: *Journal of Geophysical Research*, v. 117, B02408, doi:10.1029/2011JB008825.
- Glotzbach, C., van der Beek, P.A., and Spiegel, C., 2011, Episodic exhumation and relief growth in the Mont Blanc massif, Western Alps from numerical modelling of thermochronology data: *Earth and Planetary Science Letters*, v. 304, p. 417–430, doi:10.1016/j.epsl.2011.02.020.
- Green, P.F., 1986, On the thermo-tectonic evolution of Northern England: Evidence from fission track analysis: *Geological Magazine*, v. 123, p. 493–506, doi:10.1017/S0016756800035081.
- Green, P.F., 2002, Early Tertiary paleo-thermal effects in northern England: Reconciling results from apatite fission track analysis with geological evidence: *Tectonophysics*, v. 349, p. 131–144, doi:10.1016/S0040-1951(02)00050-1.
- Green, P.F., Duddy, I., and Bray, R.J., 1997, Variation in thermal history styles around the Irish Sea and adjacent areas: Implications for hydrocarbon occurrence and tectonic evolution, in Meadows, N.S., et al., eds., *Petroleum Geology of the Irish Sea and Adjacent Areas*: Geological Society of London Special Publication 124, p. 73–93, doi:10.1144/GSL.SP.1997.124.01.06.
- Green, P.F., Westaway, R., Manning, D.A.C., and Younger, P.L., 2012, Cenozoic cooling and denudation in the North Pennines (northern England, UK) constrained by apatite fission track analysis of cuttings from the Eastgate Borehole: *Proceedings of the Geologists' Association*, v. 123, p. 450–463, doi:10.1016/j.pgeola.2011.11.003.
- Hoke, G.D., Graber, N.R., Mescua, J.F., Giambiagi, L.B., Fitzgerald, P.G., and Metcalf, J.R., 2015, Near pure surface uplift of the Argentine Frontal Cordillera: Insights from (U-Th)/He thermochronometry and geomorphic analysis, in Sepúlveda, S.A., et al., eds., *Geodynamic Processes in the Andes of Central Chile and Argentina*: Geological Society of London Special Publication 399, p. 383–399, doi:10.1144/SP399.4.
- Holliday, D.W., 1993, Mesozoic cover over northern England: Interpretation of apatite fission track data: *Journal of the Geological Society*, v. 150, p. 657–660, doi:10.1144/gsjgs.150.4.0657.
- Jaupart, C., and Mareschal, J.C., 2005, Constraints on crustal heat production from heat flow data, in Rudnick, R.L., ed., *The Crust*: Amsterdam, Elsevier, p. 53–73.
- Majorowicz, J., and Minea, V., 2012, Geothermal energy potential in the St-Lawrence River area, Québec: *Geothermics*, v. 43, p. 25–36, doi:10.1016/j.geothermics.2012.03.002.
- Neumann, N., Sandiford, M., and Foden, J., 2000, Regional geochemistry and continental heat flow: Implications for the origin of the South Australian heat flow anomaly: *Earth and Planetary Science Letters*, v. 183, p. 107–120, doi:10.1016/S0012-821X(00)00268-5.
- Persano, C., Bishop, P., and Stuart, F.M., 2006, Apatite (U-Th)/He age constraints on the Mesozoic and Cenozoic evolution of the Bathurst region, New South Wales: Evidence for antiquity of the continental drainage divide along a passive margin: *Australian Journal of Earth Sciences*, v. 53, p. 1041–1050, doi:10.1080/08120090600923303.
- Pollack, H.N., and Cercone, K.R., 1994, Anomalous thermal maturities caused by carbonaceous sediments: *Basin Research*, v. 6, p. 47–51, doi:10.1111/j.1365-2117.1994.tb00074.x.
- Thomson, K., Underhill, J.R., Green, P.F., Bray, R.J., and Gibson, H.J., 1999, Evidence from apatite fission track analysis for the post-Devonian burial and exhumation history of the northern Highlands, Scotland: *Marine and Petroleum Geology*, v. 16, p. 27–39, doi:10.1016/S0264-8172(98)00064-6.
- Tiley, R., White, N., and Al-Kindi, S., 2004, Linking Paleogene denudation and magmatic underplating beneath the British Isles: *Geological Magazine*, v. 141, p. 345–351, doi:10.1017/S0016756804009197.
- Turcotte, D.L., and Schubert, G., 2002, *Geodynamics*: Cambridge, UK, Cambridge University Press, 456 p., doi:10.1017/CBO9780511807442.
- Wildman, M., Brown, R.W., and Gleadow, A.J.W., 2015, Post break-up tectonic inversion across the southwestern cape of South Africa: New insights from apatite and zircon fission track thermochronometry: *Tectonophysics*, v. 654, p. 30–55, doi:10.1016/j.tecto.2015.04.012.

Manuscript received 7 February 2017  
Revised manuscript received 26 April 2017  
Manuscript accepted 30 April 2017

Printed in USA