



Originally published as:

Garcin, Y., Deschamps, P., Ménot, G., de Saulieu, G., Schefuß, E., Sebag, D., Dupont, L. M., Oslisly, R., Brademann, B., Mbusnum, K. G., Onana, J.-M., Ako, A. A., Epp, L. S., Tjallingii, R., Strecker, M. R., Brauer, A., Sachse, D. (2018): Reply to Clist et al.: Human activity is the most probable trigger of the late Holocene rainforest crisis in Western Central Africa. - *Proceedings of the National Academy of Sciences of the United States of America (PNAS)*, 115, 21, pp. E4735—E4736.

DOI: <http://doi.org/10.1073/pnas.1805582115>

REPLY TO CLIST ET AL.:

Human activity is the most probable trigger of the late Holocene rainforest crisis in Western Central Africa

Yannick Garcin^{a,1}, Pierre Deschamps^b, Guillemette Ménot^c, Geoffroy de Saulieu^d, Enno Schefuß^e, David Sebag^{f,g,h}, Lydie M. Dupont^e, Richard Oslisly^{d,i}, Brian Brademann^j, Kevin G. Mbusnum^k, Jean-Michel Onana^{l,m}, Andrew A. Akoⁿ, Laura S. Epp^o, Rik Tjallingii^r, Manfred R. Strecker^a, Achim Brauerⁱ, and Dirk Sachse^p

Clist et al. (1) challenge our conclusions (2), criticizing our archaeological synthesis to maintain that the late Holocene rainforest crisis (LHRC) in Western Central Africa (WCA) was not triggered by human activity.

Clist et al. (1) claim that the archaeological ¹⁴C dates we used were not critically evaluated, as we were more concerned with the quantity of dates rather than their quality. However, a careful reading of our article (2) and its associated *SI Appendix* unambiguously documents that we cautiously weighted the radiometric dates by applying a binning in space and time to correct for investigator bias and oversampling at different sites, following refs. 3–5; this was done before combining and rigorously testing the dates.

Clist et al. (1) emphasize that the initial regional increase in human occupation in the study region lags the inception of the local LHRC at Lake Barombi by 200 y. Instead, chronological uncertainties—of ~160 y (95% confidence range) for the Lake Barombi LHRC (2) and of ~100–200 y for the population dynamics proxy used (5)—indicate that this apparent time lag is statistically insignificant, particularly as the compared sites straddle the Equator. Here seasonal interhemispheric shifts in atmospheric CO₂ cause additional uncertainties in ¹⁴C calibration (6). Conversely, the studies (7, 8) cited by Clist et al. (1) ignore the chronological uncertainties of ¹⁴C dates and compile, without rigorous statistical tests,

regional data with limited chronological control and sedimentary hiatuses, which precludes any correlation and attribution to a single event.

The signal in the Lake Barombi sediments—the best-dated sedimentary record so far in WCA—is local and strongly amplified due to the high sensitivity of this small basin to vegetation changes. The timing of the LHRC may differ from site to site, also implying chronological leads or lags with the regional trends of human occupation.

The local return to nearly “full” C₃ vegetation after the Lake Barombi LHRC (at ~2,000 cal y BP) may either reflect a regeneration of the rainforest or a replacement of C₄ crops by C₃ crops, which is compatible with the pronounced coeval increase in oil palm (a C₃ plant) use and human occupation in the region (2, 9), contradicting the claim of Clist et al. (1).

Contrary to the study (10) cited by Clist et al. (1) to suggest that charcoal is absent at Lake Barombi, our pollen analysis on core B14 includes quantifiable amounts of charcoal (see <https://doi.pangaea.de/10.1594/PANGAEA.884675>), which indicate an up to fourfold increase in charcoal at the inception of the LHRC, together with an increase in sedimentation rate.

Finally, Clist et al. (1) provide no viable alternative explanation for the observed environmental change in the Lake Barombi region at the time of the LHRC,

^aInstitute of Earth and Environmental Science, University of Potsdam, 14476 Potsdam, Germany; ^bAix Marseille Univ, CNRS, IRD, INRA, Coll France, CEREGE, 13545 Aix-en-Provence, France; ^cUniv Lyon, Ens de Lyon, Université Lyon 1, CNRS, LGL-TPE, 69342 Lyon, France; ^dPALOC, IRD, MNHN, 75005 Paris, France; ^eMARUM—Center for Marine Environmental Sciences, University of Bremen, 28359 Bremen, Germany; ^fNormandie Université, UNIROUEN, UNICAEN, M2C, CNRS, 76000 Rouen, France; ^gHSM, Univ Montpellier, CNRS, IRD, 34095 Montpellier, France; ^hInstitute of Earth Surface Dynamics, Geopolis, University of Lausanne, 1015 Lausanne, Switzerland; ⁱAgence Nationale des Parcs Nationaux, BP 20379 Libreville, Gabon; ^jSection 5.2, Climate Dynamics and Landscape Evolution, GFZ-German Research Centre for Geosciences, 14473 Potsdam, Germany; ^kAix Marseille Univ, CNRS, LCE, 13545 Aix-en-Provence, France; ^lDepartment of Plant Biology, Faculty of Sciences, University of Yaoundé I, Yaoundé, Cameroon; ^mHerbier National du Cameroun, Institut de Recherche Agricole pour le Développement, Yaoundé, Cameroon; ⁿHydrological Research Centre, Institute of Geological and Mining Research, Yaoundé, Cameroon; ^oAlfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, 14473 Potsdam, Germany; and ^pSection 5.1, Geomorphology, GFZ-German Research Centre for Geosciences, 14473 Potsdam, Germany

Author contributions: Y.G., P.D., G.M., G.d.S., E.S., D. Sebag, L.M.D., R.O., B.B., K.G.M., J.-M.O., A.A.A., L.S.E., R.T., M.R.S., A.B., and D. Sachse wrote the paper.

The authors declare no conflict of interest.

Published under the [PNAS license](#).

¹To whom correspondence should be addressed. Email: yannickgarcin@yahoo.fr.

Published online May 8, 2018.

particularly in the absence of any significant hydrological change, as we clearly demonstrate. The first widespread and intensive forest clearances at ~2,500 cal y BP in East Africa were associated with the

arrival of iron-using early farming communities (11). Similarly, our observations are compatible with the hypothesis that the LHRC in the Lake Barombi region was driven by human activity.

-
- 1 Clist B, et al. (2018) Did human activity really trigger the late Holocene rainforest crisis in Central Africa? *Proc Natl Acad Sci USA* 115:E4733–E4734.
 - 2 Garcin Y, et al. (2018) Early anthropogenic impact on Western Central African rainforests 2,600 y ago. *Proc Natl Acad Sci USA* 115:3261–3266.
 - 3 Shennan S, et al. (2013) Regional population collapse followed initial agriculture booms in mid-Holocene Europe. *Nat Commun* 4:2486.
 - 4 Timpson A, et al. (2014) Reconstructing regional population fluctuations in the European Neolithic using radiocarbon dates: A new case-study using an improved method. *J Archaeol Sci* 52:549–557.
 - 5 Crema ER, Habu J, Kobayashi K, Madella M (2016) Summed probability distribution of ¹⁴C dates suggests regional divergences in the population dynamics of the Jomon Period in Eastern Japan. *PLoS One* 11:e0154809.
 - 6 McCormac FG, et al. (2004) SHCal04 Southern Hemisphere calibration, 0–11.0 cal kyr BP. *Radiocarbon* 46:1087–1092.
 - 7 Bostoen K, et al. (2015) Middle to late Holocene Paleoclimatic change and the early Bantu expansion in the rain forests of Western Central Africa. *Curr Anthropol* 56:354–384.
 - 8 Maley J, et al. (2018) Late Holocene forest contraction and fragmentation in central Africa. *Quat Res* 89:43–59.
 - 9 Sowunmi MA (1999) The significance of the oil palm (*Elaeis guineensis* Jacq.) in the late Holocene environments of west and west central Africa: A further consideration. *Veg Hist Archaeobot* 8:199–210.
 - 10 Giresse P, Maley J, Brenac P (1994) Late Quaternary palaeoenvironments in the Lake Barombi Mbo (West Cameroon) deduced from pollen and carbon isotopes of organic matter. *Palaeogeogr Palaeoclimatol Palaeoecol* 107:65–78.
 - 11 Marchant R, et al. (2018) Drivers and trajectories of land cover change in East Africa: Human and environmental interactions from 6000 years ago to present. *Earth Sci Rev* 178:322–378.