




Originally published as:

Murray, K. E., Braun, J., Reiners, P. W. (2018): Toward Robust Interpretation of Low-Temperature Thermochronometers in Magmatic Terranes. - *Geochemistry Geophysics Geosystems (G3)*, 19, 10, pp. 3739—3763.

DOI: <http://doi.org/10.1029/2018GC007595>



Geochemistry, Geophysics, Geosystems

RESEARCH ARTICLE

10.1029/2018GC007595

Key Points:

- Intrusive magmatism may produce important first-order effects on patterns of low-temperature thermochronometer cooling
- Analytical approximations and numerical models predict cooling ages in and around crustal plutons in steadily exhuming environments
- Large plutons emplaced in the middle crust at 10–15-km depth can reset low-temperature thermochronometers in the upper crust

Correspondence to:

K. E. Murray,
murrayke@umich.edu

Citation:

Murray, K. E., Braun, J., & Reiners, P. W. (2018). Toward robust interpretation of low-temperature thermochronometers in magmatic terranes. *Geochemistry, Geophysics, Geosystems*, 19, 3739–3763. <https://doi.org/10.1029/2018GC007595>

Received 5 APR 2018

Accepted 7 SEP 2018

Accepted article online 11 SEP 2018

Published online 12 OCT 2018

Toward Robust Interpretation of Low-Temperature Thermochronometers in Magmatic Terranes

Kendra E. Murray^{1,2} , Jean Braun^{3,4,5}, and Peter W. Reiners¹ 

¹Department of Geosciences, University of Arizona, Tucson, AZ, USA, ²Now at Department of Earth and Environmental Sciences, University of Michigan–Ann Arbor, Ann Arbor, MI, USA, ³ISTerre, Université Grenoble-Alpes, CNRS, Grenoble Cedex, France, ⁴Helmholtz Centre Potsdam, GFZ German Research Center for Geosciences, Potsdam, Germany, ⁵Institute of Earth and Environmental Science, University of Potsdam, Potsdam, Germany

Abstract Many regions central to our understanding of tectonics and landscape evolution are active or ancient magmatic terranes, and robust interpretation of low-temperature thermochronologic ages in these settings requires careful attention to the drivers of rock heating and cooling, including magmatism. However, we currently lack a quantitative framework for evaluating the potential role of magmatic cooling—that is, post-magmatic thermal relaxation—in shaping cooling age patterns in regions with a history of intrusive magmatism. Here we use analytical approximations and numerical models to characterize how low-temperature thermochronometers document cooling inside and around plutons in steadily exhuming environments. Our models predict that the thermal field a pluton intrudes into, specifically the ambient temperatures relative to the closure temperature of a given thermochronometer, is as important as the pluton size and temperature in controlling the pattern and extent of thermochronometer resetting in the country rocks around a pluton. We identify one advective and several conductive timescales that govern the relationship between the crystallization and cooling ages inside a pluton. In synthetic vertical age-elevation relationships (AERs), resetting next to plutons results in changes in AER slope that could be misinterpreted as past changes in exhumation rate if the history of magmatism is not accounted for. Finally, we find that large midcrustal plutons, such as those emplaced at ~10–15-km depth, can reset the low-temperature thermochronometers far above them in the upper crust—a result with considerable consequences for thermochronology in arcs and regions with a history of magmatic activity that may not have a surface expression.

1. Introduction

Igneous rocks are commonly targeted for studies of the timing, rate, and spatial patterns of exhumation because they typically contain abundant accessory minerals useful for low-temperature thermochronology. Interpreting the thermal histories of currently or formerly magmatically active regions is, however, complicated by transient heating and cooling within and adjacent to igneous bodies; interaction between advective and conductive cooling; and transient changes in the regional geothermal gradient. These complexities are seldom quantified or discussed in studies that rely on low-temperature thermochronologic data, in part because independent constraints on past variation of the geothermal gradient are rare. Although numerical models commonly used to interpret thermochronologic data predict the widely recognized effects of topography, relief change, and/or variations in the exhumation rate on the upper crustal geothermal gradient and cooling age patterns (e.g., Braun, 2003; Ehlers, 2005a), this top-down perspective still attributes rock cooling patterns entirely to exhumation—a common assumption that is one of the most important and rarely addressed limitations of using thermochronology to document exhumation and erosion.

This is of particular concern in many regions central to our current understanding of continental tectonics. In the magmatic arcs of the American Cordillera, for example, landscape evolution studies have used low-temperature thermochronology to infer coupling between precipitation and erosion (Reiners et al., 2003); tectonic responses to Pleistocene global climate cooling modulated by glacial erosion (Thomson et al., 2010); and how the development of relief and surface elevation reflect tectonic influences on crustal buoyancy (House et al., 1998). Magmatic cooling can also impact the detrital record. Lovera et al. (1999) present a thermokinetic model and statistical framework for using detrital K-feldspar ⁴⁰Ar/³⁹Ar ages to

derive the exhumation histories of the eroded Peninsular Ranges batholith. Taking a different approach, Malusà et al. (2011) present a conceptual model for how crystallization ages (zircon U/Pb) and cooling ages (biotite K/Ar, zircon, and apatite fission track) should change with stratigraphic level in a basin that archives the unroofing of a volcanic-plutonic complex in the Alps. They report that not accounting for magmatic cooling in the detrital source region can result in apparent geological paradoxes—for example, rapid exhumation in a source terrane contemporaneous with slow sediment accumulation rates in the basin—because thermal relaxation can be misinterpreted as rapid exhumation.

At best, studies of landscape evolution in magmatic terranes use available high-temperature geochronology from locally exposed plutons to argue that magma emplacement happened long enough before exhumation that the cooling ages are unrelated to magmatic thermal relaxation (e.g., Guillaume et al., 2013; Murray et al., 2016; Reiners et al., 2003). However, using the lag time between high-temperature crystallization ages and low-temperature cooling ages to make this case is only valid when there is a framework for identifying what a sufficient lag time is—a framework we currently lack. One objective of our study is to quantitatively evaluate this approach.

Early thermochronology studies discussed the cooling of upper crustal plutons as a function of both magmatic thermal relaxation and exhumation (e.g., Harrison & Clarke, 1979a) because higher temperature methods—and questions—dominated thermochronology at that time. Harrison and Clarke (1979a) were the first to model the cooling of a pluton while accounting for geothermal flux, rock uplift, and radioactive heat production. This offered a quantitative check that the proposed post-magmatic history (Harrison & Clarke, 1979b) is generally valid, but it explicitly did not seek a best-fit thermal history. Another classic study (Calk & Naeser, 1973) sampled granitic country rocks immediately adjacent to a small basaltic plug to validate fission-track annealing kinetics in accessory minerals.

A few studies have explicitly modeled the thermal effects of upper crustal magmatism on bedrock thermochronometers from regions with a known history of magmatism. Low-temperature thermochronologic ages locally reset by plutonism are useful for identifying hydrothermal ore deposits (Fu et al., 2010; McInnes et al., 2005). In a study of the timing of Rio Grande rift footwall tilting and exhumation, House et al. (2003) evaluate the role of hypothesized midcrustal plutonism on apatite cooling ages on the rift flanks. They conclude that there is little geologic evidence for a pluton of the size and position required to generate the observed age patterns and therefore attribute the cooling to exhumation. Regional magmatism in South Africa's Karoo basin is thought to be coeval with Gondwana breakup, so Brown et al. (1994) use simple thermal models of various magmatic processes to quantitatively predict the possible effects on the Karoo sedimentary units, concluding that additional processes like magmatic hydrothermal activity likely played a substantial role. In each case, whether magmatic cooling was part of the preferred cooling age interpretation, the potential effects of magmatism on the thermal history of the upper crust were clear and substantial.

Spatial and temporal changes in regional heat flow—commonly attributed to lithosphere-scale processes—have also been interpreted from low-temperature thermochronologic data sets. For example, cooling age patterns have been attributed to the thermal effects—mostly surface uplift and erosion—of transient slab windows (Guenther et al., 2010; Guillaume et al., 2013) and hotspots (Taylor & Fitzgerald, 2011). Alternatively, flat slabs may refrigerate the lithosphere (e.g., the Laramide slab in the Sierra Nevada; Dumitru et al., 1991). These studies present conceptual models of these processes, but with the exception of Roy et al. (2004), they do not quantitatively investigate the thermal effects of the proposed geodynamic processes. Moreover, they do not consider magma migration as an important mechanism for rapid and efficient heat transfer in the upper crust.

In sum, despite the clear sensitivity of low-temperature thermochronometers to nonexhumational cooling (Ehlers, 2005b), few studies—especially those relying on the lowest-temperature apatite thermochronometers—address this potential complication when interpreting cooling ages.

This study is motivated by the need to broadly quantify how the thermal effects of plutonism manifest in the low-temperature thermochronologic record. We use analytical and numerical thermal models (Figure 1) to predict temperatures and cooling age patterns around and inside thermal anomalies in the upper crust. Our approach is grounded in Laplace's analytical approximation (Figure 1a) to one-dimensional (1D) unsteady heat conduction problem in an infinite region (Turcotte & Schubert, 2002), which finds the general shape and pattern of heating around an instantaneously emplaced heat source and is useful for predicting




Figure 1. Summary of this study's modeling approaches. (a) Laplace's solution to one-dimensional unsteady heat conduction problem in an infinite region provides simple predictions of the temperature and cooling age patterns around plutons. (b) The numerical model Pecube solves for the thermal field for different exhumation rates and calculates cooling ages in and around a pluton. The simplest model in Pecube is similar to Laplace's solution in three-dimensions (3D), with a tall narrow pluton emplaced in the upper 10 km of the crust. (c) Small tabular plutons and (d) large tabular plutons have more geologically realistic shapes in 3D. (e) Models with tabular plutons emplaced at 12–15-km depths explore the effects of midcrustal magmatism on upper crustal thermochronometers.

Table 1
Abbreviations and Units in This Study

| Symbol | Unit | Description |
|----------------------------------------------------------------------------|--------------------|------------------------------------------------------------------------------------------------------------------------|
| T_Z | °C or K | Ambient temperature of the country rocks at depth Z at the time of magmatic activity |
| T_C | °C or K | Nominal closure temperature of a thermochronometer |
| Z_C | km | Depth of T_C |
| T_P | °C or K | Temperature of the igneous body at instantaneous emplacement |
| T_B | °C or K | Temperature at the base of a model |
| PRZ | — | Partial-retention zone; a range of temperatures (or depths) where He is neither fully retained nor lost from a crystal |
| 1D approximation from Laplace's solution (after Turcotte & Schubert, 2002) | | |
| y | km | Distance in 1D from center of half space |
| T_{\max} | K | Peak temperature at position y |
| t | s | Time |
| t_{\max} | s | Time after instantaneous emplacement T_{\max} is reached at position y |
| Q | J/km ² | Heat content of pluton per unit area of pluton-country rock interface |
| ρ | kg/km ³ | Rock density |
| c | kJ/kg K | Heat capacity |
| κ | km ² /s | Thermal diffusivity |
| b | km | Pluton half width |
| l | kJ/kg | Latent heat of crystallization |
| Pecube model results | | |
| τ | Ma | Cooling age |
| τ_L | Ma | Cooling age in the landscape undisturbed by magmatic activity, i.e., the <i>landscape</i> or <i>background</i> age |
| τ_X | Ma | Pluton crystallization age, taken to be the time since <i>emplacement</i> in the model |
| ξ | — | Normalized cooling age (τ/τ_L) |

the general patterns of cooling age resetting. We also use a modified version of the three-dimensional (3D) numerical model Pecube (Braun, 2003; Braun et al., 2011) to solve the heat advection-conduction equation and calculate cooling ages in simple emplacement-exhumation scenarios. This provides a guide for further investigations of the geologic circumstances where magmatic heating and cooling should be considered during cooling age interpretation.

2. Methodology

2.1. Low-Temperature Thermochronometric Systems

Low-temperature thermochronometers are the only tools available for documenting the thermal history of the upper crust on 10⁶- to 10⁸-year timescales. The commonly used (U-Th)/He and fission-track systems in apatite and zircon rely on the time-dependent radioactive decay of U and Th nuclides and the temperature-dependent retention and/or annealing of their daughter products (⁴He and fission tracks) in individual crystals. On geologic timescales, the kinetics of the diffusion or annealing of these radiogenic daughter products are commonly described as a thermochronometric system's closure temperature, T_C (Dodson, 1973), which is a function of the cooling rate. The (U-Th)/He chronometers in apatite (AHe) and zircon (ZHe) document rock cooling below ~40–80 and ~140–220 °C, respectively. The fission track chronometers in these minerals (AFT and ZFT) document rock cooling below ~90–120 and ~220–260 °C, respectively.

2.2. Modeling

We use two approaches for modeling the thermal effects of magmatic heating and cooling on thermochronometers (see Table 1 for a summary of abbreviations and model variables). First, we use an analytical approximation of Laplace's solution to 1D unsteady heat conduction problem in an infinite region (Turcotte & Schubert, 2002) to make simple predictions about the patterns of heating around a pluton instantaneously emplaced into a region of the crust of specific ambient temperature. Then, we use the thermokinematic model Pecube (Braun, 2003; Braun et al., 2011) to capture how exhumation rate affects the pattern and

Table 2
Thermokinematic Parameters Used in Pecube Models

| Parameter | Value | Unit |
|--------------------------------------------------|--------|----------------------|
| Crustal thickness | Varies | km |
| Thermal diffusivity | 25 | km ² /Myr |
| Basal crustal temperature (T_B) ^a | Varies | °C |
| Sea-level temperature | 0 | °C |
| Atmospheric lapse rate | 0 | °C/km |
| Crustal heat production | 0 | °C/Myr |
| Igneous body temperature (T_P) | 750 | °C |

Note. Elastic parameters not reported because isostatic response to relief change is not modeled.

^aModel basal temperature selected assuming a 25 °C/km linear geothermal gradient and a given crustal thickness.

magnitude of thermochronometer resetting in and around plutons. Pecube is an established tool for quantitative interpretation of thermochronologic data. It solves the 3D heat transport equation in a crustal block experiencing rock uplift and surface erosion beneath a time-evolving surface topography. Pecube computes the time-temperature histories of particles that move through the calculated thermal structure and computes the resulting cooling ages for a range of thermochronometers. The modifications we made for this study are among many in the continuous development of Pecube as its applications have widened to address how thermochronometers document tectonic and geomorphic processes (Braun et al., 2011).

In Pecube, we simulate the thermal effects of pluton emplacement by holding a region of the model domain at a constant high temperature for a finite time before letting it cool. In these models, the hot region is a

rectangular prism or cube. Relevant parameters include the hot region's holding temperature (°C), its depth (km), its half width, half length, and half height (km), as well as the heating start and stop time (Ma). Between the heating start and stop times, model nodes in the prescribed region are isothermally held. To capture the transient thermal response during this heating event and immediately after, the model time step is adapted to be shorter than the conductive timescale of pluton cooling set by the pluton size.

Other than these modifications, the code is as described in Braun (2003) and Braun et al. (2011). The model geometry represents a crustal block. Its dimensions are scaled such that the x-y dimensions are ≥ 10 times the pluton half-width b_{XY} to ensure that the result is not influenced by the assumed no-flux boundary conditions on the lateral boundaries. We impose a constant basal temperature that is obtained by assuming a linear 25 °C/km geothermal gradient and a given crustal thickness. The initial temperature distribution is obtained by running the model to conductive-advective steady state (given the exhumation rate) prior to pluton emplacement. We also assume a flat topography that does not evolve with time, so we can focus our analysis on the effect of the thermal perturbation caused by the pluton alone. In order to observe the bedrock cooling age patterns, after pluton emplacement we impose a total exhumation of 9 or 10 km in each Pecube simulation, which exhumes all rocks that were shallower than or close to the closure depths of the highest-temperature chronometer we model (the zircon He system) at the time of magmatic activity. The length of each model run, therefore, scales with the exhumation rate. We stack the exhumed material into synthetic cross sections to visualize the results. Pecube model parameters are given in Table 2.

As in many thermochronologic studies, the closure temperature (T_C) as defined by Dodson (1973) provides a useful framework here. It is, however, a mathematical description of thermochronometer behavior for monotonic cooling histories that has limited quantitative meaning in rocks reheated by magmatic activity. The use of fixed T_C in the 1D model results presented here is, therefore, a simplification of thermochronometer behavior. Pecube, however, does not rely on the T_C concept. Instead, it solves the full production-diffusion (or -annealing) equations to calculate cooling ages given the model-predicted rock thermal histories. We report closure depths (Z_C) in Pecube models set by the imposed constant exhumation rate.

2.3. Definitions

We use *pluton* as a general term to describe a volume of igneous material but do not intend to imply a particular emplacement mechanism, fraction of melt, or bulk composition. However, as described above, the pluton temperatures we use are most relevant for granitic compositions. We suggest that the cooling age patterns discussed here can be associated with the entire range of igneous bodies, from sills to batholiths. *Country rocks* are the material outside the pluton, which, for simplicity, we assume have the same physical properties (i.e., thermal diffusivity and density) as the plutonic material.

We present model results as patterns of temperature, T , or thermochronologic cooling age, τ , inside and outside the modeled plutons. To compare the model results for different exhumation rates, we compare cooling ages to two other ages that are measurable in the model and, potentially, in nature: (1) the cooling age the rocks would have if not perturbed by magmatism (i.e., the cooling age of country rocks in the landscape far

from the pluton; τ_L) and (2) the crystallization age of the pluton (i.e., the pluton's geochronologic age; τ_X), which is defined in our models as the time elapsed since the end of isothermal heating.

To examine the pattern of cooling ages affected by magmatic cooling, we normalize τ by τ_L (for the same thermochronometric system) to get a value ξ . Where $\xi = 1$, cooling ages are unaffected by magmatic cooling and $\tau = \tau_L$. Where $\xi < 1$, the cooling ages τ are younger than τ_L and to some extent document the effect of heating by the pluton. In the country rocks outside of and adjacent to the pluton, we define the *resetting aureole* to be the part of the thermal aureole where $\xi < 1$ for a particular thermochronometric system. We note that in most thermochronologic data sets, intrasample cooling age reproducibility is commonly $\sim 10\%$, and therefore in nature, only ξ values of < 0.9 would be a significant thermochronologic signal.

2.4. Simplifying Magma Body Energetics

Our modeling approach reduces the complex and poorly constrained process of magma emplacement to a simple thermal event. We do not attempt to mimic thermal pulses from individual emplacement events that likely build intrusive complexes and their source regions (Annen, 2011; Annen et al., 2015; Glazner et al., 2004), nor do we account for the dependence of thermal diffusivity on temperature (Nabelek et al., 2012), the effects of magma or fluid convection within or surrounding the intrusion (Norton & Knight, 1977; Norton & Taylor, 1979; Parmentier & Schedl, 1981), or a particular style of magma emplacement. Although important at some spatial and temporal scales (Jaeger, 1964), these factors present substantial uncertainties and numerical challenges that are beyond the scope of this study. Moreover, at the $> 10^6$ -year timescales that thermochronometers document, the thermal effects of many of these processes would effectively smooth out, so our simple models should capture the first-order effects of intrusive systems on the thermal architecture of the upper crust.

Additionally, we are only modeling the pluton's sensible heat and therefore do not directly account for the latent heat of crystallization, which constitutes a considerable fraction of a cooling pluton's total energy. Because the sensible heat is proportional to the temperature difference between the pluton and the country rocks, the latent heat is a larger fraction of the total heat available to raise the temperature of the country rocks in more deeply emplaced magma bodies. For example, for a granitic magma at 750°C , latent heat is $\sim 25\%$ of the total if emplaced into 50°C country rocks but $\sim 70\%$ if the country rocks are 650°C .

In models without a complete mathematical treatment of magma energetics, like ours, it is common to account for latent heat by simply increasing the initial pluton temperature (Calk & Naeser, 1973; Harrison & Clarke, 1979a). However, the actual temperatures experienced by the country rocks, and not just the total pluton heat, are important for thermochronology. So instead, in our Pecube models, we hold a pluton at a constant temperature of 750°C for thousands of years during *emplacement*. This mimics how latent heat would function as a thermal buffer (Huber et al., 2009) that keeps the pluton hotter for longer than sensible heat alone. The emplacement duration time, along with the pluton and country rock temperatures, controls how much heat is added and therefore has a substantial effect on the extent of thermochronometer resetting. Therefore, we use a conservatively cold pluton temperature (750°C) because it is below the solidus for most magma compositions. Incrementally constructed magma bodies cool below the magma solidus between injection events in all but rare high-flux scenarios (Annen, 2011). In most simulations, we scale the holding time to the pluton volume because the time-averaged rate of pluton construction is broadly correlated to total volume (de Saint-Blanquat et al., 2011). We use a rate of $10^{-2} \text{ km}^3/\text{Myr}$, slightly faster than the average observed in compilations of data from a diverse range of igneous body size, composition, and tectonic setting (10^{-1} to $10^{-4} \text{ km}^3/\text{Myr}$; de Saint-Blanquat et al., 2011).

Although choosing a pluton temperature of 750°C is meant to simulate the thermal effects of granitic magmas, the difference between the latent and sensible heat outlined above provides a useful scaling relationship relating intrusions of varying temperature (e.g., granite and basalt). To generate the same amount of heat as a granite magma at shallow crustal levels (i.e., country rocks at 25°C), $\sim 33\%$ less basalt volume is required; for deeper crustal levels (country rocks at 650°C), $\sim 60\%$ less is required.

3. Advection and Conduction Timescales

Our purpose is to investigate the thermal perturbation caused by the emplacement of a pluton and how it affects the distribution of thermochronometer ages otherwise set by exhumation-driven cooling. For

simplicity, we assume a constant exhumation rate in each of our models. The cooling history of a rock inside or in the vicinity of the pluton is governed by the competition between advective (or exhumation-driven) heat transport, conductive cooling through the Earth's surface, and conductive cooling of the pluton. We can measure the efficiency of each process through their respective characteristic timescale. The timescale for conductive cooling of the crust is given by

$$\varphi_C = \frac{L^2}{\kappa} \quad (1)$$

and the timescale for heat advection by

$$\varphi_A = \frac{L}{\dot{E}} \quad (2)$$

where L is a length scale (km; usually the thickness of the crust or of the crustal layer being exhumed), \dot{E} is the exhumation rate (km/Myr), and κ is the thermal diffusivity (km²/Myr). The relative efficiency of these two processes can be assessed by the ratio of these two timescales, which defines the so-called Péclet number:

$$Pe = \frac{\varphi_C}{\varphi_A} = \frac{\dot{E}L}{\kappa} \quad (3)$$

When $Pe \gg 1$, advective transport dominates and there is a strongly depth-dependent geothermal gradient. When $Pe \ll 1$, conductive cooling dominates and the geothermal gradient is quasi-uniform with depth. The timescale for conductive cooling of the pluton is, in turn, given by

$$\varphi_C^{\text{pluton}} \propto \frac{b^2}{\kappa} \quad (4)$$

where b is the half width (or radius) of the pluton. The ratio of the pluton conductive timescale to the advection timescale measures how rapidly the pluton cools while being exhumed:

$$Pe^{\text{pluton}} = \frac{\varphi_C^{\text{pluton}}}{\varphi_A} = \frac{\dot{E}b^2}{L\kappa} \quad (5)$$

The competition between the advection and conduction of heat also dictates what part of the rock cooling history (magmatic or exhumational) a thermochronometer documents, given that \dot{E} sets the advection timescale (equation (2)) and the pluton size (a length scale) sets a conductive timescale proportional to its half-width, b (equation (4)).

When a 3D pluton is emplaced into a block with a geothermal gradient, two different pluton dimensions are important length scales (Appendix A). The conductive cooling timescale at the center of the pluton is set by the pluton thickness in the Z dimension, because the thickness is parallel to the principal direction of heat conduction toward the cold surface. The cooling timescale in the resetting aureole next to the pluton is set by the pluton's X and Y dimensions. Therefore, in our 3D models, we distinguish between the half width in the Z dimension (b_z) and the half widths in the X and Y dimensions, which are equal (b_{xy}).

4. 1D Analytical Approximation

4.1. Laplace's Solution

In country rocks adjacent to a pluton, the width of a resetting aureole is a function of the ambient temperature at the depth, T_z , and the closure temperature of the thermochronologic system, T_c . An approximate expression for the width of the resetting aureole as a function of T_z and T_c can be derived from the Laplace's solution to 1D unsteady heat conduction problem in an infinite region (Turcotte & Schubert, 2002; equation (4)–(157)) given by

$$T = \frac{1}{2\sqrt{\pi\kappa t}} \int_{-\infty}^{\infty} \bar{T}(y') \exp\left[-\frac{(y-y')^2}{4\kappa t}\right] dy' \quad (6)$$

where t is time, c is heat capacity, ρ is density, κ is thermal diffusivity, y is a horizontal spatial coordinate, and $\bar{T}(y)$ is the initial temperature anomaly from a heat source (Table 1). If we only solve for the temperature in the resetting aureole far from the pluton, $y \gg b_{XY}$, where b_{XY} is the pluton half width, we can approximate the pluton as a planar source of heat located at $y = 0$ (Figure 1a). The solution gives the temperature-time evolution at a distance y from an instantaneously emplaced heat source:

$$T(y, t) = T_Z + \frac{Q}{2pc\sqrt{\pi\kappa t}} \exp\left[-\frac{y^2}{4\kappa t}\right] \quad (7)$$

where Q is the amount of heat contributed by the pluton:

$$Q = \rho[c(T_P - T_Z) + l]2b_{XY} \quad (8)$$

T_P is the pluton temperature, and l is the latent heat of crystallization (which here we assume is 0). From equation (7), we can derive the time at which rocks at position y reach their maximum temperature (t_{\max}):

$$t_{\max}(y) = \frac{y^2}{2\kappa} \quad (9)$$

The peak temperature at that time is

$$T_{\max}(y) = T_Z + \frac{Q}{\rho c y} \left(\frac{1}{2\pi e}\right)^{0.5} \quad (10)$$

By combining equations (8) and (10), we estimate the distance y where T_{\max} equals a temperature of interest, for example, the T_C of a thermochronologic system:

$$y_{\max} \cong \left(\frac{T_P - T_Z}{T_C - T_Z}\right) \left(\frac{b_{XY}}{2}\right) \quad (11)$$

Rocks within y_{\max} distance of the pluton experienced peak temperatures greater than T_C (Figure 2a).

This approximation highlights a principal feature of resetting aureoles: Their width increases exponentially with T_Z , as T_Z approaches the T_C of the thermochronologic system of interest (Figure 2a).

4.2. Schematic Cooling Age Patterns

Laplace's simple solution highlights two spatial relationships that control whether a country rock's temperature exceeds a thermochronometer's T_C as a result of localized heating next to a pluton: (1) the rock's depth or, more exactly, its ambient temperature T_Z at the time of magmatism, and (2) its proximity to the pluton. All else being equal, these two variables control the shape of the temperature field and therefore the spatial extent of the resetting aureole (Figures 2a and 3a). If we assume a pluton was emplaced into a crustal block where the cooling age pattern prior to magmatism was set by steady and spatially uniform exhumation (e.g., Figure 3b) and that this steady exhumation continued long after magmatism, we can make simple schematic predictions of the horizontal and vertical cooling age patterns that plutonism should produce.

Along horizontal transects (Figure 2a) where $T_Z < T_C$, the cooling ages range from $\tau = \tau_X$ close to the pluton to $\tau \cong \tau_L$ far from the pluton, and the width of the thermochronometer resetting aureole is a function of the T_Z at the time of magmatism (Figure 2b). At depths where T_Z is closer to T_C , the resetting aureole is wider, and under our assumption of steady-exhumation, the difference between reset and unreset ages is also smaller (Figure 2b). This is because at the time of magmatism, the cooling ages in rocks where T_Z is close to T_C are very close to zero (Figure 3b) and so resetting results in a smaller change in age than in rocks where $T_Z \ll T_C$. As time passes and these transects progressively exhume, the pluton age τ_X gets older and the difference between τ_X and τ_L decreases until the rocks being exhumed were at $T_Z > T_C$ at the time of plutonism, had a cooling age of $\tau = 0$ at that time, and were therefore unperturbed by pluton heating.

In vertical transects (Figure 3), a pluton's perturbation of the existing age- T_Z relationship (Figure 3b) is a function of the horizontal distance from the pluton (Figure 3c). The vertical thickness of the resetting aureole is




Figure 2. We use Laplace's solution to find the shape of the resetting aureole for a single thermochronometric system and the resulting schematic horizontal cooling age patterns. (a) Cross section through a thermochronometer resetting aureole was constructed by solving equation (10) for a range of ambient temperatures. The width of the resetting aureole is a function of temperature at depth Z , T_z . The closer T_z is to the closure temperature T_c , the wider the resetting aureole. Horizontal lines labeled 1–4 indicate the depth of cooling age transects in panel (b) at the time of magmatism. (b) Schematic horizontal cooling age transects given the dimensions of the resetting aureole in (a). Close to the pluton, the cooling ages are completely reset and are set by the pluton emplacement (τ_X). Far from the pluton, the cooling ages are unaffected by magmatic heating and are set by the exhumation rate (τ_L). Note that as time passes and deeper rocks are exhumed, the pluton's crystallization age τ_X gets older. In rocks at shallow depths at the time of magmatism (i.e., at cold T_z ; transect 1), the resetting aureole is narrow, and the age difference between reset and unreset ages is large. Transects exhumed later through the same surface from increasingly greater depths and hotter T_z have increasingly wider resetting aureoles but smaller cooling age differences across the transect.




Figure 3. Laplace's solution predicts the spatial relationship between thermochronometric systems with different closure temperatures and suggests vertical cooling age patterns at different distances from the pluton for a single system. (a) Analytical approximations of the resetting aureoles of the four commonly used low-temperature thermochronometers, created by solving equation (10) with the following parameters: closure temperature T_C for each system given a $10^\circ\text{C}/\text{Myr}$ cooling rate (Reiners & Brandon, 2006), a pluton with half-width $b = 0.5$ km and temperature $T_P = 750^\circ\text{C}$, and ambient temperatures T_Z from 10 to 250°C . Vertical lines indicate the position of vertical profiles in panels (c) and (d). (b) Zircon He ages as a function of T_Z for a constant $10^\circ\text{C}/\text{Myr}$ cooling rate (i.e., a constant exhumation rate \dot{E}) as predicted by the thermal history modeling code HeFTy (Ketchum, 2005) using standard zircon He diffusion kinetics. At the surface, the cooling age is τ_L . Where $T_Z > T_C$, the cooling age is zero. (c) Zircon He age versus T_Z patterns (i.e., age-depth or age-elevation relationships) immediately after pluton cooling at three locations: the edge of the pluton (square), at $y/b = 3$ (diamond) and $y/b = 10$ (circle) distance from the pluton center. For the purposes of generally describing the cooling age patterns that would result from resetting around plutons using Laplace's approximation, we assume that when the peak $T > T_C$, a chronometer is completely reset. A heating event would perturb the age- T_Z relationship in panel (b), creating sharp changes in the slope of a vertical profile at the transitions into an out of a resetting aureole. (d) As constant (exhumation-driven) cooling at a rate of $10^\circ\text{C}/\text{Myr}$ continues after pluton emplacement, the perturbed age- T_Z relationship is progressively exhumed. At this rate it takes ~ 20 Myr to exhume all the rocks that were colder than the zircon He T_C ($\sim 183^\circ\text{C}$) at the time of pluton emplacement and entirely re-establish the age- T_Z relationship in (b). Rocks from depths where $T_Z \gg T_C$ at the time of magmatism do not document magmatic cooling because they had a zero age at the time of pluton emplacement. Each thermochronologic system would have these patterns, which would be nested as a function of T_C .

greatest in transects closest to the pluton. This resetting extends up from rocks where $T_Z = T_C$ at the time of plutonism. The resetting aureole in a vertical transect manifests as significant kinks in the age- T_Z relationship, which is progressively exhumed (Figure 3d) during post-magmatic exhumation.

Several key insights follow from this simple 1D treatment. First, the ambient temperature T_Z is as important as the pluton size and temperature in controlling the amount of extra heat added to the crust by a pluton and therefore its thermal effect (equation (7)). Second, resetting aureoles for low-temperature thermochronometers with different T_C will be nested. The higher T_C systems will have narrower aureoles that extend to higher T_Z than those with lower T_C (Figure 3a). Third, in a steadily exhuming landscape with no additional perturbations, if a pluton's crystallization age τ_X is older than the cooling age of the country rocks τ_L , then the magnitude of exhumation since pluton emplacement is greater than the depth to the closure

temperature, Z_C , of the thermochronometric system (final panels of Figures 2b and 3d). Finally, in vertical profiles, as would be obtained in nature from a borehole or vertical sampling transect, the resetting aureole manifests as significant changes in the slope of the age- T_Z (i.e., age-depth and age-elevation) relationship (Figures 3c and 3d). Such trends in low-temperature thermochronologic data are commonly used to infer past changes in erosion rate, but here they reflect a history of steady erosion during which exhumation-driven cooling was perturbed by a transient thermal event.

Laplace's approximation has limited utility beyond broadly characterizing the features to expect from thermochronometer resetting around plutons, because it does not include the kinetics that govern how low-temperature thermochronometers respond to temperature change over time. For example, a $T \geq T_C$ condition must be long lived in order to fully reset a thermochronometer, and therefore, the location of $T_{\max} = T_C$ (Figure 3a) overestimates the width of thermochronometer resetting aureoles. Next, we use a numerical model that solves the thermochronologic age equations to more robustly explore how low-temperature thermochronometers document interactions between magmatic and exhumation-driven cooling.

5. Upper-Crustal Plutons

Using Pecube, we model several pluton geometries and emplacement depths to examine the effects of heating from a single pluton in the upper crust (Figures 1b–1d). We compare cooling age patterns from models where the exhumation rate is set at three different but constant values of $\dot{E} = 0.01, 0.1$, or 1 km/Ma. We limit the results presented here to the apatite and zircon He systems, because they span a representative temperature sensitivity range for low-temperature thermochronometers.

5.1. Simplest 3D Models

We first model a tall, narrow pluton (Figure 1b) that resembles the infinite-sheet heat source of Laplace's solution. The crustal block (model domain) is 30×30 km² by 10 km thick and has a pluton of dimensions $b_{XY} = 6$ km and $b_Z = 8$ km kept at 750 °C for 10 kyr.

5.1.1. Horizontal Transects

Horizontal cooling age patterns (Figure 4) follow the predictions from Laplace's solution (Figure 2b). Inside and immediately adjacent to the pluton, the cooling ages are approximately equal to the pluton crystallization age ($\tau \cong \tau_X$). Moving away from the pluton in each transect where $T_Z < T_C$, we predict the existence of a partially reset zone, where cooling ages are older than the pluton but younger than the landscape cooling age ($\tau_X < \tau < \tau_L$). Farther from the pluton, magmatic cooling did not affect cooling ages, so $\tau = \tau_L$ and $\xi = 1$.

In order to examine the effects of different exhumation rates, \dot{E} , on the cooling age patterns predicted inside and around plutons for two different thermochronometric systems, we have to select comparable horizontal transects from each model (Figure 4). For this, we have to find comparable depths relative to the thermochronometer T_C (i.e., Figure 2). Simply normalizing T_Z by T_C does not account for the possibility of a nonlinear geothermal gradient in scenarios with rapid \dot{E} and large Pe numbers (equation (3); Figure 5) and the resulting dependence on the depth of the closure temperature, Z_C , on \dot{E} . The temperature at a depth z for a given Pe value is

$$T(z) = T_B \frac{1 - \exp\left[-\frac{Pe z}{B}\right]}{1 - \exp[-Pe]}. \quad (12)$$

and the depth to the closure temperature is

$$z_c = -\frac{B}{Pe} \ln\left(1 - \frac{T_c(1 - \exp[-Pe])}{T_B}\right) \quad (13)$$

where B is the thickness of the model in the z direction and T_B is the temperature at the base of the model. We compare transects from the same Z/Z_C (i.e., depth normalized to Z_C) depths:

$$\frac{Z}{Z_C} = \frac{\ln\left(1 - \frac{T_Z(1 - \exp[-Pe])}{T_B}\right)}{\ln\left(1 - \frac{T_C(1 - \exp[-Pe])}{T_B}\right)}. \quad (14)$$




Figure 4. Horizontal normalized cooling age ($\xi = \tau/\tau_L$) transects from normalized Z/Z_C depths (a) Transects are across the simplest pluton exhumed at $\dot{E} = 0.01$ (b, c), 0.1 (d, e), and 1 km/Myr (f, g). The gray field indicates rocks in pluton domain held at 750°C . Where $\xi = 1$, the cooling age documents only exhumation-driven cooling. In horizontal transects from slowly exhuming scenarios and from rocks that were cold relative to the chronometer T_C , the normalized cooling age ξ in the pluton is approximately equal to the Z/Z_C , and the transition between reset to unreset ages is sharp. Transects from rocks close to Z_C at the time of magmatism and from rapidly exhuming models deviate from this $\xi = Z/Z_C$ trend and have wider resetting aureoles.




Figure 5. Temperature-depth profiles from a one-dimensional numerical model that solves the vertical heat conduction-advection equation. Depth and temperature are normalized by the thickness of the layer being exhumed and the maximum (fixed) temperature at the base of the layer, respectively. Each panel (a, b, c) corresponds to different values of the exhumation velocity that translates into different values of the Peclet number, Pe . The thick blue line is the steady-state temperature profile set by \dot{E} . We assume that a pluton has been emplaced at $t = 0$ in the center of the layer. It has a thickness of 0.2 and an initial temperature of 1.5 (i.e., 1.5 times the temperature at the base of the layer), which creates the initial temperature profile (orange line). Black lines are the temperature-depth profiles at 10 time steps separated by the time it takes to advect a point by 0.05 , such that at the end of the 10 time steps, a point that was located in the center of the layer at $t = 0$ (shown by the red circles) reaches the surface. We see that the temperature anomaly caused by the pluton decays much more rapidly in the case of a small value of Pe ($Pe = 0.1$) compared to a large value of Pe ($Pe = 10$). This is simply because in the case of fast exhumation (large Pe), the time necessary to advect the rock toward the surface is much shorter, but the conductive cooling time is unchanged, and as a result, the conductive relaxation of the temperature anomaly caused by the pluton is less efficient. This implies that the rock being exhumed at the center of the pluton remains hotter, that is, above the initial, steady-state temperature profile (thick gray line), and potentially above the closure temperature of the thermochronometer under consideration, for a longer period of time, therefore resulting in a smaller value of ξ (τ/τ_L) at a given Z/Z_C depth.

The transects of normalized cooling age ξ as a function of y/b_{xy} from depths where $Z/Z_C = 0.25, 0.5, 0.75, 1$, and 1.25 (Figure 4a) show the progressive widening of the resetting aureole with increasing Z/Z_C (Figure 2b). The maximum width of the resetting aureoles around these plutons is substantially smaller than that suggested by Laplace's solution, because fully or partially resetting a thermochronometer requires significant time at temperatures at or exceeding T_C , not just instantaneously reaching a peak temperature greater than the T_C set.

As shown in Figure 4, faster \dot{E} results in wider resetting aureoles. At slow rates (Figures 4b–4e), the resetting aureoles for both thermochronologic systems are at most $\sim b_{xy}$ (3 km) wide, with slightly wider aureoles at 0.1 km/Myr. When $\dot{E} = 1$ km/Myr (Figures 4f and 4g), the resetting aureole extends as much as $2b_{xy}$ from the pluton edge.

Within the pluton, comparing a transect's Z/Z_C value to the ξ value (τ/τ_L) reveals the effect of \dot{E} on pluton cooling and thereby the cooling ages of plutonic rocks. At shallow depths and in slowly exhumed models (Figures 4b–4e), the Z/Z_C value is approximately equal to ξ . For example, in the model with $\dot{E} = 0.01$ km/Myr (Figures 4b and 4c), the transect from a shallow depth where $Z/Z_C = 0.25$ has pluton cooling ages that are 25% of the landscape's background cooling age (i.e., $\xi = 0.25$) and therefore $\xi = Z/Z_C$. This reflects very rapid cooling to below the thermochronometer closure temperatures after emplacement. In deeper transects very close to the closure depth ($Z/Z_C \sim 1$; Figure 4), the plutonic rocks are hotter than T_C for longer because little additional heat is required to maintain $T > T_C$. As a result, the pluton cooling age is younger than the cooling age we would predict from rapid postemplacement cooling to ambient temperatures. Therefore, the normalized age is less than the normalized depth ($\xi < Z/Z_C$). In models with rapid \dot{E} , this effect is magnified and $\xi < Z/Z_C$ at much shallower depths (Figures 4f and 4g). The deviation from $\xi \approx Z/Z_C$ is greater for the apatite He results than for the zircon He results (Figures 4f and 4g).

5.1.2. Advection and Pluton Cooling

Delayed cooling in advection-dominated systems with rapid \dot{E} is predicted by these 3D models. It manifests as a deviation from $Z/Z_C \approx \xi$ inside a pluton (section 5.1.1 and Figures 4f and 4g) and reflects how advection controls the efficiency of dissipating a transient heat anomaly in the crust.

If the preintrusion geothermal gradient is approximately linear because exhumation rates are slow (i.e., $\dot{E} \ll 1$), the Z/Z_C value is also the ratio between the pluton age and the landscape cooling age (τ_X/τ_L). Because we defined $\xi = \tau/\tau_L$, the ratio between ξ and Z/Z_C is also the ratio between the pluton's cooling age and emplacement age:

$$\frac{\xi}{Z/Z_C} = \frac{\tau}{\tau_X} \quad (15)$$

Plutonic rocks that rapidly cooled after emplacement to the ambient T set by \dot{E} have $\tau \cong \tau_X$ and $\xi/(Z/Z_C) = 1$, which is why $\xi \approx Z/Z_C$ inside the pluton and parts of the resetting aureoles in the slowly exhumed and shallowest transects in Figure 4. If cooling is delayed, then the lag time between the pluton crystallization and cooling is longer, $\tau < \tau_X$, and therefore $\xi < Z/Z_C$.

We use a simpler 1D conduction-advection model to illustrate this behavior. In Figure 5, we compute the shape of a geothermal gradient (in normalized depth vs. temperature space) set by three different exhumation rates that correspond to Pe values of 0.1, 1, and 10. We perturb these temperature-depth gradients with the same temperature anomaly and predict temperature-depth profiles at 10 subsequent time steps evenly spaced over the time it takes to advect a point from the center of the model—inside the pluton—to the surface. The amount of time this advection takes scales directly with the Pe value; it takes 100 times longer for 10 time steps to elapse in the $Pe = 0.1$ model than the $Pe = 10$ model. In contrast, the time it takes for conductive relaxation of the pluton is constant. As a result, in slowly advecting models, the system conductively dissipates the heat anomaly introduced by the pluton much more rapidly than it advects it (Figure 5a). In rapidly advecting models, rocks from a given depth take much longer to cool relative to their exhumation time to the surface (Figure 5c). They potentially remain hotter than T_C longer and have smaller ξ (τ/τ_L) values.

For lower-temperature thermochronometers, it takes less time for rocks to get from Z_C to the surface (i.e., the advection timescale is shorter), and the system has less time to conductively dissipate the pluton heat over that time. This explains why thermochronometers with colder closure temperatures and shallower Z_C (i.e., the apatite He system here; Figures 4b, 4d, and 4f) are more perturbed ($\xi < Z/Z_C$) at rapid \dot{E} .

5.1.3. Vertical Profiles

Examining these results along a vertical cross section (Figure 6) offers a perspective relevant to sampling strategies common in low-temperature thermochronology. Contoured normalized cooling ages ($\bar{\epsilon}$) from a cross section through the model (Figure 6, left panels) show resetting aureoles widening as Z approaches Z_C and the significantly wider resetting aureoles for plutons emplaced in rapidly exhuming scenarios. The difference between τ and τ_X in vertical profiles at different positions relative to the pluton (Figure 6, middle panels) shows the magnitude of the resetting in Myr. To demonstrate how these age patterns would manifest in a vertical transect, we construct synthetic age-elevation relationships (AERs; Figure 6, right panels). The AERs are for a snapshot in time at the end of the model run, when 9 km of exhumation has occurred since pluton emplacement. It takes two orders of magnitude longer to exhume this thickness of crust when $\dot{E} = 0.01$ km/Myr than when $\dot{E} = 1$ km/Myr, so the cooling and crystallization ages also vary by orders of magnitude between the models.

As predicted by the Laplace solution (Figure 3), AERs vary as a function of distance from the pluton (Figure 6, right panels). In the plutonic rock at the center of the model, the AER is subvertical at shallow levels where $\tau \cong \tau_X$ because $T_Z \ll T_C$ at the time of magmatism and the thermochronometers are documenting rapid post-magmatic thermal relaxation. In the plutonic rock emplaced near $T_Z = T_C$, the lag time between τ and τ_X is greater and the AER shallows, approaching a slope defined by \dot{E} . In the country rocks, an AER slope can be shallow or steep compared to the slope set by \dot{E} and can have three kinks in slope. This is captured best by the zircon He system (Figure 6b, right panels, green, purple, and yellow curves). At shallow levels, the AER slope reflects the background \dot{E} because the rocks are unreset. At hotter T_Z , the AER slope shallows substantially with increasing partial resetting, then it steepens to subvertical where it is fully reset by the pluton heating, and then finally it shallows again at $T_Z \gg T_C$ to reflect the constant \dot{E} .

5.1.4. Consequences for Age-Elevation Interpretations

Slope changes in vertical profile AERs are commonly interpreted as past changes in exhumation rate. AER slopes in resetting aureoles (Figure 6) could be wrongly interpreted as past variability in \dot{E} if not recognized as resulting from magmatic thermal perturbations. In most settings, a vertical crustal section available for sampling is no more than 3–4 km thick. To explore how this may manifest in nature, we calculated the apparent exhumation rate over a 1-km moving average using cooling ages along each AER to examine the predicted magnitude of this effect inside the pluton and in the country rocks.


Inside the pluton at shallow levels, the subvertical AER reflects rapid thermal relaxation and if interpreted as \dot{E} would overestimate the rate by an order of magnitude or more; greater overestimation would occur in slowly exhuming landscapes. At depths near Z_C , the AER shallows and approaches a slope that reflects the actual \dot{E} , so the \dot{E} overestimation diminishes in rocks from depths where T_Z was close to T_C . At faster \dot{E} , the AER slope inside the pluton does not reflect the actual \dot{E} until rocks were nearly 1 km deeper than the Z_C at the time of magmatism, although \dot{E} is only overestimated by a factor of 2 in rocks below the Z_C in these scenarios. These trends in the plutonic rocks are simple compared to those in the resetting aureoles because τ cannot be older than τ_X .

The AERs within the country rock resetting aureoles change significantly in slope and could result in either overestimation or underestimation of \dot{E} . Consider a transect at distance $1.67b_{xy}$ from the pluton center (Figure 6b, purple vertical profiles). At shallow levels, the rocks are unreset, and the slope of the AER reflects the actual \dot{E} . In rocks slightly deeper and partially reset, the AER is much shallower and would underestimate the \dot{E} significantly or suggest a fossil partial-retention zone (PRZ). Slightly deeper, the rocks are completely reset by magmatism and follow the same pattern as the cooling ages in the plutonic rocks. AERs closer to the pluton have the most extreme variability in slope, and a greater proportion of their vertical profile is dominated by ages fully or partially reset by magmatism.

5.2. Tabular Plutons

Geophysical and field-based studies of upper crustal plutons suggest that many are tabular (Annen et al., 2015; Cruden & McCaffrey, 2001). The tabular plutons in our models have two different sizes but have equal height to length ratios, and the amount of time they are held at 750 °C is scaled by volume assuming a time-averaged emplacement rate of 10^{-2} km³/Myr (de Saint-Blanquat et al., 2011) rounded to the nearest thousand years. The small pluton is 3 km in diameter and 1 km thick (Figure 1c), with a volume

(A) APATITE HE RESULTS



(B) ZIRCON HE RESULTS





Figure 6. Pecube model results from the simplest pluton scenarios (Figure 1b) stacked into vertical cross sections and synthetic age-elevation relationships (AERs). (a) Apatite He ages predicted from models with exhumation rates, \dot{E} , of 0.01, 0.1, and 1 km/Myr. Left panels present normalized cooling age ($\xi = \tau/\tau_L$) contoured in vertical cross section. Where $\xi < 1$, red-yellow colors indicate the contrast between the cooling age, τ , and the unreset landscape age, τ_L , in the resetting aureole. Dotted line indicates the depth of the T_C set by \dot{E} . As predicted by the one-dimensional solution, the resetting aureole is widest in rocks that were close to the Z_C at the time of magmatism. Vertical colored lines indicate the location of profiles shown in the middle and right panels. These vertical profiles report the age difference (in Myr) between the reset and unreset ages (middle panels) and the AER shape (right panels) inside the pluton and at different distances away. Note that the sampling resolution is every 500 m and starting at 1-km depth, so some details of the apatite He patterns are lost. (b) Zircon He ages predicted from the same models.

APATITE HE RESULTS



ZIRCON HE RESULTS




Figure 7. Model predictions for small tabular plutons (Figure 1c), shown as contoured ξ (τ/τ_L) values in vertical cross section as in Figure 6.

of 9 km^3 and a heating time of 1 kyr. The large pluton is 15 km in diameter and 5 km thick (Figure 1d), with a volume of $125,000 \text{ km}^3$ and a heating time of 113 kyr. We place them at very shallow crustal levels (pluton roof at 2-km depth) or slightly deeper (pluton roof at 5-km depth) and exhume them at 0.1 and 1-km/Myr rates.

Unlike the simplest pluton modeled in section 5.1, the small tabular plutons modeled here do not occupy a significant fraction of the crustal column, so their emplacement depth relative to the chronometer Z_C is the principal variable controlling the pattern and magnitude of the magmatic cooling documented by that thermochronometer (Figure 7). When emplaced shallowly (Figures 7a and 7b), the small tabular plutons overlap with the apatite He Z_C , so the resetting aureole widens with proximity to Z_C and is wider in the more rapidly exhuming scenario. In contrast, the same pluton emplaced 3 km deeper (Figures 7c and 7d) has no measurable effect on the cooling ages and $\xi = 1$ through the entire crustal section with the exception of a narrow band of ages $\xi \approx 0.9$ near the Z_C in the rapidly exhuming model (Figure 7c).

The small shallow plutons are emplaced several kilometers above the zircon He Z_C (Figures 7e and 7f); therefore, the resetting aureole is very narrow and is widest at the middle of the pluton. The deeper tabular plutons sit just above the zircon He Z_C , so the resetting aureole is wider and widens moderately with depth, but there is only modest contrast ($\xi > 0.75$) between the cooling ages inside and outside the resetting aureole (Figures 7g and 7h).

The large shallow pluton (Figure 8) intersects the apatite He Z_C like its smaller counterpart and generates a proportionally wider resetting aureole (Figures 8a and 8b). The large tabular pluton emplaced well below the apatite He Z_C generates significant resetting in the country rocks more than 3 km above it (Figures 8c and 8d). This effect is much greater in the model with $\dot{E} = 1 \text{ km/Myr}$ (Figure 8c). The large tabular plutons emplaced at both depths intersect the zircon He Z_C (Figures 8e–8h). As in the apatite He system, the zircon He ages kilometers above the pluton roof are reset

by this large pluton (Figures 8g and 8h).

The timescale of conductive cooling is mostly controlled by pluton thickness (equation (4), Appendix A) and is therefore longer for the larger pluton. This, along with the longer heating time, is the principal reason the thermal footprint of these two plutons is different. Additionally, when \dot{E} is faster, a larger part of the crustal column cools during the thermal relaxation of the pluton and captures part of that magmatic cooling. This follows from the scaling relationships described in section 3.

In contrast, the extensive resetting in the country rocks above a large pluton, in some cases kilometers above the pluton roof (Figure 8), is a new insight from the thermokinematic modeling. It suggests that plutons emplaced deep in the crust could cause a significant thermal perturbation in the rocks above them. We use models of midcrustal plutons to further characterize this phenomenon and its potential implications.

6. Midcrustal Plutons Reset Upper-Crustal Country Rocks

6.1. 3D Numerical Models

To explore how low-temperature thermochronometric cooling ages document thermal perturbations from plutons emplaced far below thermochronometer closure depths, we examine cooling age patterns from the upper 10 km of the crust from Pecube models with midcrustal plutons (Figure 1e). These plutons are tabular, $16 \text{ km}^2 \times 6 \text{ km}$ thick and emplaced at two different midcrustal depths: 12 and 15 km. These depths correspond to the brittle-ductile transition in the crust, where we assume magmatic ascent is interrupted (Burov




Figure 8. Model predictions for large tabular plutons (Figure 1d), shown as contoured ξ (τ/τ_L) values in vertical cross section as in Figure 6.

et al., 2003). We hold them at 750 °C for 1 or 10 Myr, to simulate a large plutonic complex constructed over 1 Myr or a midcrustal magma body acting as a recharging magma reservoir, respectively. The plutons are exhumed toward the surface at 0.1 km/Myr, so 100 m and 1 km of rock exhumation, respectively, occur while they are held at 750 °C.

In all cases, the zircon and apatite He chronometers in rocks far above the midcrustal plutons and close to Z_c depths at the time of magmatism are significantly reset (Figure 9). The size of the upper-crustal resetting aureole scales with the proximity of the pluton and how long it was hot. But in most cases, this midcrustal heating results in kilometer-scale thick sections of crust with cooling ages approximately equal to the timing of magmatic activity ($\tau \sim \tau_X$). Additionally, AERs are characterized by perturbations similar to those in country rocks immediately next to shallower upper-crustal plutons (Figure 6). These cooling age patterns could be interpreted as a change in exhumation rate, a period of rapid erosion, or a fossil PRZ. Instead, they are simply the result of a transient increase in the geothermal gradient due to a thermal perturbation at depth.

6.2. 1D Analytical Approximation

To generalize the consequences of midcrustal plutonism for low-temperature thermochronometers, we return to Laplace's solution to the 1D unsteady heat conduction problem (equation (5)) but apply it now along the vertical coordinate, z . We use this solution to determine the extent of the thermal perturbation in the upper crust caused by a midcrustal pluton by assuming the pluton is much wider than it is thick (Appendix B). The maximum temperature reached at a depth z following the rapid cooling of a pluton with half-thickness b_z buried at a depth z_0 is given by

$$T_{\max}(z) = \frac{\Delta T_0 b_z}{z_0 - z} \sqrt{\frac{1}{2\pi e}} + T_0 \frac{z}{z_0}. \quad (16)$$

T_0 is the temperature at depth Z_0 before the pluton emplacement and $T_0 + \Delta T_0$ is the initial temperature of the pluton. We compute the depth, z_1 , where the temperature will exceed a given closure temperature, T_c , at any time after the instantaneous cooling of the pluton:




Figure 9. Large tabular plutons emplaced in the middle crust (Figure 1e) heat the rocks above in the upper crust and reset the apatite and zircon He systems. The exhumation rate $\dot{\epsilon}$ is 0.1 km/Myr and sets the apatite He τ_L (25 Ma) and the zircon He τ_L (68 Ma). Resetting is shown in contoured ξ (τ/τ_L) values as well as the difference (Myr) between the landscape age τ_L and age τ (in Myr) at three positions: in the middle of the model directly above the center of the pluton, above the edge of the pluton, and in the rocks b_{XY} distance from the pluton edge. Age differences of $<10\%$ between τ_L and τ (i.e., $\xi > 0.9$) are not mapped in the ξ plot but are evident in the age difference versus depth plot. Synthetic age-elevation relationships (AERs) at each of these positions were constructed assuming 10 km of exhumation (and 100 Myr elapsed) since pluton emplacement ended. (a) Tabular pluton emplaced at 15-km depth and hot for 1 Myr. Rocks in the crust above this pluton close to Z_C at the time of magmatism are fully or partially reset by this thermal event. (b) Pluton at 12-km depth, hot for 1 Myr. (c) Pluton at 15-km depth, hot for 10 Myr. (d) Pluton at 12-km depth, hot for 10 Myr.

$$z_1 = z_0 \frac{T_0 + T_c - \sqrt{(T_0 + T_c)^2 - 4(T_0 T_c - T_0 \Delta T_0 b_Z / z_0 \sqrt{2\pi e})}}{2T_0}. \quad (17)$$

We also compute the maximum temperature that would be reached at any depth if the pluton was held at a constant temperature for a very long (infinite) time:

$$T_{\max}^{\infty} = (T_0 + \Delta T_0) \frac{z}{z_0} \quad (18)$$

and from it, we can determine the depth, z_2 , where the temperature will exceed the same closure temperature, T_c :

$$z_2 = \frac{z_0 T_c}{T_0 + \Delta T_0} \quad (19)$$

In Figure 10, we show how these two depths z_1 and z_2 vary as a function of T_c and compare them to the unperturbed closure depth Z_C .

This simple analytical prediction agrees with the numerical model predictions in 3D (Figure 9). It provides a general guide for the potential extent of low-temperature thermochronometer resetting in the upper crust above large long-lived or short-lived magmatic bodies (of specific size) emplaced at depths much greater than Z_C .




Figure 10. Laplace's analytical solution used to determine the crustal depths that would experience $T > T_c$ as the result of the presence of an instantaneously cooled (blue field) or long-lived (infinite, red field) pluton located directly below in the middle crust. The black line is the depth versus temperature in an unperturbed linear geothermal gradient of 30 °C/km. This line is also the depth of any temperature of interest, such as a T_c . Blue and red lines identify the shallower depths where the same temperature would be (e.g., where $T = T_c$) as a result of the instantaneously cooled or long-lived pluton, respectively. The pluton temperature ($T_0 + \Delta T_0$) is 750 °C. Broad ranges for the temperature sensitivities of the three lowest-temperature thermochronometric systems identified for reference. This approach provides a general guide for the extent of resetting above midcrustal plutonic systems of different sizes and depths, for example, (a) $b_z = 2.5$ km, $Z_0 = 15$ km, and (b) $b_z = 1$ km, $Z_0 = 10$ km.

7. Implications for Low-Temperature Cooling Age Interpretations

Heating from intrusive magmatism most extensively resets low-temperature thermochronometers in country rocks that are colder than, but close to, the chronometer closure temperature (i.e., in the PRZ or partial-annealing zone) at the time of magmatic activity. Around plutons emplaced at or above thermochronometer Z_c depths, the resetting zones are localized aureoles around the pluton (Figures 6, 7, and 8). Our models predict that midcrustal magmatism can also reset thermochronometers at PRZ depths—thermochronometers in rocks that are kilometers above the pluton itself (Figures 8d and 9). It is useful to describe these resetting extents as *local* and *regional* effects, respectively.

Regional resetting may be especially relevant for felsic magmatic systems, which require million-year-scale crustal residence times to produce evolved magma compositions if they were originally fed by mafic magmas. Additionally, because isolated, shallowly emplaced plutons are merely the top of a more voluminous magmatic system that plumbs the middle and lower crust, in some regions it may be critical to consider both the local heating around observed individual upper-crustal plutons and the regional thermal effects of the related magmatic system at depth. Such systems would have an additive heating effect that is thought to produce low-pressure metamorphic belts (Barton & Hanson, 1989; Hanson & Barton, 1989).

It appears that the common practice of using the crystallization ages of magmatic rocks mapped at the surface as a spatial-temporal guide for interpreting cooling ages can be insufficient in regions with a history of midcrustal magmatism. Model results predict that plutons emplaced in the middle crust and not yet exhumed are capable of resetting low-temperature thermochronometers in the upper crust without any surface expression. In other words, the low-temperature thermochronologic record of a midcrustal pluton's emplacement forms at shallower depths than, and will be exhumed before, the pluton itself. Such regional resetting may be particularly important in magmatic arcs, where plutonism occurs continuously at geologic timescales, and therefore surface mapping can document only part of the magmatic history relevant to thermochronologic interpretations.

The resetting signal produced by midcrustal plutons is only present in rocks that were at PRZ depths at the time of plutonic activity. Therefore, this reset zone must be subsequently exhumed to the surface (or sampled in a borehole) to be observed. For example, if it is independently known that rocks at the surface today were at apatite He partial-retention depths (i.e., ~2 km; Figure 10a) in the past at a time of known magmatic activity, the apatite He cooling ages of these rocks should be interpreted with caution. In contrast, zircon He ages from the same rocks were likely too shallow and therefore too cold to be reset

by the same magmatic event, and the pre-magmatic cooling history would be preserved in the zircon He system. For the regional magmatic cooling signal to be observed in zircon He ages at the surface today, post-magmatic exhumation of ~4–5 km (Figure 10a) would be required to exhume rocks from zircon He PRZ depths.

This study simulates bedrock cooling ages but has implications for detrital thermochronology. Sedimentary sections with material shed during the unroofing of magmatic terranes can have detrital cooling ages that reflect both magmatic and exhumational cooling of the source region. When shallowly emplaced plutons are exhumed in a source terrane, rocks from both the pluton and resetting aureole may erode and be deposited at the same time. Therefore, as in the bedrock scenarios modeled here, the high-temperature geochronometer (crystallization) ages in detrital minerals can provide an important reference age to compare to the low-temperature cooling ages (e.g., Malusà et al., 2011). In contrast, the resetting aureoles formed above midcrustal plutons will exhume before the plutons and produce a detrital thermochronologic signal that could, as in interpretations of bedrock AERs, appear to document a period of rapid exhumation in the source terrane.

This is one of many common complexities in nature that we do not directly explore here because we limited our models to steadily exhuming bedrock landscapes with no topography. Our results do, however, provide a general framework for interpreting cooling ages in magmatic terranes by clarifying the role of a region's pre- and post-magmatic history.

The pre-magmatic history sets both the cooling-age field and thermal field into which a pluton intrudes. The cooling-age field controls the age contrast between the reset, partially reset, and unreset thermochronometers (i.e., ζ). It therefore governs how magmatic cooling ages will be in manifest in the geologic record. The pre-magmatic thermal field controls the spatial extent and pattern of thermochronometer resetting at depth during magmatism. A terrane's postmagmatic exhumation history controls whether rocks that document magmatic cooling are in an accessible part of the geologic record, that is, at the surface today or incorporated into the detrital thermochronologic record.

The shape of the surface boundary condition (i.e., the premagmatic and syn-magmatic surface topography) is an important variable—especially for the lowest- T chronometers—that we do not explore here. If, for example, plutonism occurs in a region with kilometer-scale topographic relief, then plutons will intrude into a thermal field with corresponding relief in the upper crustal isotherms. Therefore, the resetting aureoles will be shaped by the topography of the surface boundary condition. Particularly interesting complications may arise if the surface boundary conditions change as a result of ongoing plutonism. For example, shallowly emplaced tabular plutons (e.g., laccoliths) commonly uplift the country rocks, and potentially the surface, above them. This creates surface topography and the potential for magmatism-driven exhumation (Gilbert, 1877). Given the number of possible variables (the thermochronometers of interest; the pluton size, depth, and emplacement time; the initial topography and its evolution over time; the timescale and spatial scale of the erosional response to uplift, etc.), it is not possible to generalize about the effects of this on the low-temperature thermochronologic record. Our modified version of Pecube is, however, capable of exploring some specific scenarios or situations applicable to a given case study.

8. Conclusions

The analytical and numerical modeling results presented here offer a new perspective on how to best interpret low-temperature thermochronologic cooling ages inside and around plutonic rocks. Laplace's analytical solution to the unsteady heat conduction problem in a 1D infinite region predicts the importance of the ambient temperature at depth, T_z , in controlling the spatial extent of thermochronometer resetting. The resetting aureole for a thermochronometric system is widest at depths close to the closure temperature T_C (Figure 2a), and therefore, the extent of resetting around a pluton is not just a function of its size and temperature. The thermokinematic model Pecube, which numerically solves the 3D heat transport equation and calculates thermochronometer cooling ages, supports the predictions from Laplace's solution and predicts how the thermal perturbations produced by simple plutons of various sizes and emplacement depths affect thermochronometers, given different constant exhumation rates.

The scaling relationships that define the Péclet number, Pe , describe how the cooling history of rocks inside and around a pluton is governed by the competition between exhumation-driven (or advective) cooling and




Figure A1. Setup for the numerical model in Appendix A.

conductive cooling of the pluton, and conductive cooling through the Earth's surface (Figure 5). In models with rapid exhumation, advective heat transport dominates. As a result, a larger section of the crust is affected by the thermal perturbation introduced by the pluton in horizontal and vertical dimensions, but the absolute difference between reset and unreset cooling ages (Myr) is small because rock cooling is very rapid and all cooling ages are very young (Figure 6).

In all cases where a pluton is emplaced at depths close to or above a thermochronometer closure temperature, the resetting aureole(s) in the surrounding country rocks creates substantial changes in the slope of vertical AERs (Figure 6). Such changes in AER slope are commonly used to infer past changes in exhumation rate or periods of rapid erosion, but here they simply reflect the magmatic perturbation of the geothermal gradient in a steadily exhuming landscape. Interpreting these AER patterns as changes in exhumation rate would result in overestimating or underestimating \dot{E} by orders of magnitude.

Our models predict that similar AER patterns can form in rocks in the upper crust directly above large mid-crustal plutons emplaced at 10–15-km depth (Figure 9). The extent of the thermal perturbation caused by these deeply emplaced plutons scales with their thickness, depth of emplacement, and longevity (Figure 10; Appendix B). The effects are focused in rocks at ambient temperatures close to T_c at the time of magmatic activity, so in order to see this effect in the geologic record, the rocks close to T_c depths at the time of regional magmatism must be subsequently exhumed. Additionally, the rocks reset by mid-crustal magmatism can be kilometers above a mid-crustal pluton, and, if so, they will exhume before the pluton itself. Therefore, plutons locally exposed in a study area may not provide a sufficient record of the magmatic cooling that may be part of a region's thermochronologic history.

The flat, constantly exhuming magmatic terranes modeled here are simple but instructive models of regions where low-temperature thermochronology is used to investigate landscape evolution. These models illustrate that (1) a region's pre-magmatic and syn-magmatic exhumation history sets the thermal and cooling-age fields into which a pluton intrudes and (2) a region's post-magmatic exhumation history controls whether rocks that document magmatic cooling are observed in the bedrock or detrital thermochronologic record. Many natural settings have topographic relief and variable exhumation that would add additional complexity. These complexities make it even more critical to recognize the first-order control the pre- and post-magmatic histories have in shaping the pattern and preservation of magmatic cooling signals in the thermochronologic record.

Appendix A: Pluton Cooling Timescales

The cooling of a pluton of thickness b_z and radius b_{xy} , emplaced at a depth z_0 in a lithosphere of thickness L (Figure A1), is characterized by a range of conductive timescales, each corresponding to a particular length scale. Assuming that the pluton thickness is smaller than its depth ($b_z < z_0$) and that its radius is larger than its thickness ($b_z < b_{xy}$), they are as follows: the timescale for cooling of the entire lithosphere (assuming that the temperature at the base of the lithosphere is constant),

$$\bar{\tau} \approx \frac{L^2}{\kappa\pi^2} \quad (\text{A1})$$

the timescale for cooling at the center of the pluton,

$$\tau_c \approx \frac{b_z^2}{\kappa\pi^2} \quad (\text{A2})$$

the timescale for cooling of the crust adjacent to the pluton but at the same depth,

$$\tau_R \approx \frac{b_{xy}^2}{\kappa\pi^2} \quad (\text{A3})$$




Figure A2. Numerical model results testing the validity of the scaling relationships described in Appendix B. The dashed lines correspond to simple exponential decay curves with a timescale equal to τ_R , τ_b , $\bar{\tau}$, and τ_0 , respectively.

and the timescale for cooling near the surface,

$$\tau_0 \approx \frac{z_0^2}{\kappa \pi^2}. \quad (\text{A4})$$

We note that each timescale is defined by the length scale defined in the direction of heat conductive transport for the location under consideration.

We tested the validity of these relationships by numerically solving the heat conduction equation in cylindrical coordinates (r, ϕ, z) , positioning the pluton along the $r = 0$ axis and assuming azimuthal symmetry. We




Figure A3. Setup for the model of midcrustal pluton in Appendix B.

included a uniform radiogenic heat production. We tracked the mean temperature, the temperature at the center of the pluton, the temperature near the surface, and the temperature next to the pluton at the same depth. We used a second-order accurate, centered difference scheme to estimate the Laplacian and an explicit scheme for the time evolution. In Figure A2, we show, for example, the time evolution of the temperature at the center of the pluton for various values of b_{XY} (Figure A2a), b_Z (Figure A2b), L (Figure A2c), and z_0 (Figure A2d). We see that the response time for the cooling at the center of the pluton is principally controlled by b_Z , the pluton thickness, as long as it is substantially smaller than its radius and its depth of emplacement.

Appendix B: Thermal Perturbation of a Midcrustal Pluton

We can use Laplace's solution to the 1D unsteady heat conduction equation (equation (5)) to determine the extent of the thermal perturbation caused by a deep crustal pluton, if we assume that the pluton is much wider than thick (i.e., it is a tabular pluton; Figure A3).

Note that Laplace's solution does not fully apply to the problem at hand, as it assumes that the medium in which the pluton intrudes is infinite. The exact solution that takes into account that the surface is characterized by a finite (nil) temperature is given by the following infinite series:

$$\Delta T(z, t) = \sum_{n=1}^{\infty} \frac{4\Delta T_0}{n\pi} \sin \frac{n\pi z_0}{L} \sin \frac{n\pi b}{L} \sin \frac{n\pi z}{L} \exp \left[-\frac{n^2 \pi^2 t \kappa}{L^2} \right]. \quad (\text{A5})$$

In Figure A4, we compare Laplace's solution to the maximum temperature predicted by the exact solution and note that they are identical in the vicinity of the pluton and only diverge by less than 10% near the surface. Because of its simplicity, we use Laplace's solution.




Figure A4. Laplace's analytical solution and the actual solution for a finite region are comparable except very close to the surface.

Acknowledgments

The code for Pecube can be found in Braun et al. (2011). This paper is theoretical, and no data are used. K. E. M. acknowledges support from a NSF GRFP award, ARCS Phoenix Chapter Prentice Scholarship, a GSA student research grant, and a P.E.O. Scholar Award. J. B. acknowledges support from GFZ German Research Center for Geosciences. Thank you to Thibault Simon-Labric for discussions on pluton heating effects on thermochronology; to Mihai Ducea, Jay Quade, and George Davis for comments on early versions of this work; and to graduate students and faculty at ISTERre for welcoming K. E. M. during visits to Grenoble, France. Additional thanks to Kiriaki Xiluri-Lauria for technical support. This manuscript benefitted from thoughtful reviews by Julie Fosdick and one anonymous reviewer.

References

- Annen, C. (2011). Implications of incremental emplacement of magma bodies for magma differentiation, thermal aureole dimensions and plutonism–volcanism relationships. *Tectonophysics*, 500(1–4), 3–10. <https://doi.org/10.1016/j.tecto.2009.04.010>
- Annen, C., Blundy, J. D., Leuthold, J., & Sparks, R. S. J. (2015). Construction and evolution of igneous bodies: Towards an integrated perspective of crustal magmatism. *Lithos*, 230, 206–221. <https://doi.org/10.1016/j.lithos.2015.05.008>
- Barton, M. D., & Hanson, R. B. (1989). Magmatism and the development of low-pressure metamorphic belts: Implications from the western United States and thermal modeling. *Geological Society of America Bulletin*, 101(8), 1051–1065. [https://doi.org/10.1130/0016-7606\(1989\)101<1051:MADOL>2.3.CO;2](https://doi.org/10.1130/0016-7606(1989)101<1051:MADOL>2.3.CO;2)
- Braun, J. (2003). Pecube: A new finite-element code to solve the 3D heat transport equation including the effects of a time-varying, finite amplitude surface topography. *Computers & Geosciences*, 29(6), 787–794. [https://doi.org/10.1016/S0098-3004\(03\)00052-9](https://doi.org/10.1016/S0098-3004(03)00052-9)
- Braun, J., van der Beek, P., Valla, P., Robert, X., Herman, F., Glotzbach, C., et al. (2011). Quantifying rates of landscape evolution and tectonic processes by thermochronology and numerical modeling of crustal heat transport using PECUBE. *Tectonophysics*, 524–525, 1–28. <https://doi.org/10.1016/j.tecto.2011.12.035>
- Brown, R., Gallagher, K., & Duane, M. (1994). A quantitative assessment of the effects of magmatism on the thermal history of the Karoo sedimentary sequence. *Journal of African Earth Sciences*, 18(3), 227–243. [https://doi.org/10.1016/0899-5362\(94\)90007-8](https://doi.org/10.1016/0899-5362(94)90007-8)
- Burov, E., Jaupart, C., & Guillou-Frottier, L. (2003). Ascent and emplacement of buoyant magma bodies in brittle-ductile upper crust. *Journal of Geophysical Research*, 108(B4), 2177. <https://doi.org/10.1029/2002JB001904>
- Calk, L. C., & Naeser, C. W. (1973). The thermal effect of a basalt intrusion on fission tracks in quartz monzonite. *The Journal of Geology*, 81(2), 189–198. <https://doi.org/10.1086/627834>
- Cruden, A. R., & McCaffrey, K. (2001). Growth of plutons by floor subsidence: Implications for rates of emplacement, intrusion spacing and melt-extraction mechanisms. *Physics and Chemistry of the Earth Part A-Solid Earth and Geodesy*, 26(4–5), 303–315. [https://doi.org/10.1016/S1464-1895\(01\)00060-6](https://doi.org/10.1016/S1464-1895(01)00060-6)
- de Saint-Blanquat, M., Horsman, E., Habert, G., Morgan, S., Vanderhaeghe, O., Law, R., & Tikoff, B. (2011). Multiscale magmatic cyclicality, duration of pluton construction, and the paradoxical relationship between tectonism and plutonism in continental arcs. *Tectonophysics*, 500(1–4), 20–33. <https://doi.org/10.1016/j.tecto.2009.12.009>
- Dodson, M. (1973). Closure temperature in cooling geochronological and petrological systems. *Contributions to Mineralogy and Petrology*, 40(3), 259–274. <https://doi.org/10.1007/BF00373790>
- Dumitru, T., Gans, P., Foster, D., & Miller, E. (1991). Refrigeration of the western. *Cordilleran lithosphere during Laramide shallow-angle subduction*, 19(11), 1145–1148. [https://doi.org/10.1130/0091-7613\(1991\)019<1145:ROTWCL>2.3.CO;2](https://doi.org/10.1130/0091-7613(1991)019<1145:ROTWCL>2.3.CO;2)
- Ehlers, T. A. (2005a). Computational tools for low-temperature thermochronometer interpretation. *Reviews in Mineralogy and Geochemistry*, 58(1), 589–622. <https://doi.org/10.2138/rmg.2005.58.22>
- Ehlers, T. A. (2005b). Crustal thermal processes and the interpretation of thermochronometer data. *Reviews in Mineralogy and Geochemistry*, 58(1), 315–350. <https://doi.org/10.2138/rmg.2005.58.12>
- Fu, F. Q., McInnes, B. I. A., Evans, N. J., & Davies, P. J. (2010). Numerical modeling of magmatic–hydrothermal systems constrained by U–Th–Pb–He time–temperature histories. *Journal of Geochemical Exploration*, 106(1–3), 90–109. <https://doi.org/10.1016/j.jgexplo.2009.09.001>
- Gilbert, G. K. (1877). *Report on the geology of the Henry Mountains*. Washington, DC: U.S. Government Printing Office.
- Glazner, A., Bartley, J., Coleman, D., Gray, W., & Taylor, R. (2004). Are plutons assembled over millions of years by amalgamation from small magma chambers? *GSA Today*, 14(4), 4–12. [https://doi.org/10.1130/1052-5173\(2004\)014<0004:APAOMO>2.0.CO;2](https://doi.org/10.1130/1052-5173(2004)014<0004:APAOMO>2.0.CO;2)
- Guenther, W. R., Barbeau, D. L. Jr., Reiners, P. W., & Thomson, S. N. (2010). Slab window migration and terrane accretion preserved by low-temperature thermochronology of a magmatic arc, northern Antarctic Peninsula. *Geochemistry Geophysics Geosystems*, 11, Q03001. <https://doi.org/10.1029/2009GC002765>
- Guillaume, B., Gautheron, C., Simon-Labric, T., Martinod, J., Roddaz, M., & Douville, E. (2013). Dynamic topography control on Patagonian relief evolution as inferred from low temperature thermochronology. *Earth and Planetary Science Letters*, 364, 157–167. <https://doi.org/10.1016/j.epsl.2012.12.036>
- Hanson, R. B., & Barton, M. D. (1989). Thermal development of low-pressure metamorphic belts: Results from two-dimensional numerical models. *Journal of Geophysical Research*, 94(B8), 10,363–10,377. <https://doi.org/10.1029/JB094iB08p10363>
- Harrison, T., & Clarke, G. K. (1979b). Model of the thermal effects of igneous intrusion and uplift as applied to Quottoon pluton, British Columbia. *Canadian Journal of Earth Sciences*, 16(3), 411–420. <https://doi.org/10.1139/e79-039>
- Harrison, T. M., & Clarke, G. K. C. (1979a). A model of the thermal effects of igneous intrusion and uplift as applied to Quottoon pluton, British Columbia. *Canadian Journal of Earth Sciences*, 16(3), 411–420. <https://doi.org/10.1139/e79-039>
- House, M. A., Kelley, S. A., & Roy, M. (2003). Refining the footwall cooling history of a rift flank uplift, Rio Grande rift, New Mexico. *Tectonics*, 22(5), 1060. <https://doi.org/10.1029/2002TC001418>
- House, M. A., Wernicke, B. P., & Farley, K. A. (1998). Dating topography of the Sierra Nevada, California, using apatite (U–Th)/He ages. *Nature*, 396(6706), 66–69. <https://doi.org/10.1038/23926>
- Huber, C., Bachmann, O., & Manga, M. (2009). Homogenization processes in silicic magma chambers by stirring and mushification (latent heat buffering). *Earth and Planetary Science Letters*, 283(1–4), 38–47. <https://doi.org/10.1016/j.epsl.2009.03.029>
- Jaeger, J. C. (1964). Thermal effects of intrusions. *Reviews of Geophysics*, 2(3), 443–466. <https://doi.org/10.1029/RG002i003p00443>
- Ketcham, R. A. (2005). Forward and inverse modeling of low-temperature thermochronometry data. *Reviews in Mineralogy and Geochemistry*, 58(1), 275–314. <https://doi.org/10.2138/rmg.2005.58.11>
- Lovera, O., Grove, M., Kimbrough, D., & Abbott, P. (1999). A method for evaluating basement exhumation histories from closure age distributions of detrital minerals. *Journal of Geophysical Research*, 104(B12), 29,419–29,438. <https://doi.org/10.1029/1999JB900082>
- Malusà, M. G., Villa, I. M., Vezzoli, G., & Garzanti, E. (2011). Detrital geochronology of unroofing magmatic complexes and the slow erosion of Oligocene volcanoes in the Alps. *Earth and Planetary Science Letters*, 301(1–2), 324–336. <https://doi.org/10.1016/j.epsl.2010.11.019>
- McInnes, B., Evans, N., Fu, F., & Garwin, S. (2005). Application of thermochronology to hydrothermal ore deposits. *Reviews in Mineralogy and Geochemistry*, 58(1), 467. <https://doi.org/10.2138/rmg.2005.58.18-498>
- Murray, K. E., Reiners, P. W., & Thomson, S. N. (2016). Rapid Pliocene–Pleistocene erosion of the central Colorado Plateau documented by apatite thermochronology from the Henry Mountains. *Geology*, 44(6), 483–486. <https://doi.org/10.1130/G37733.1>
- Nabelek, P. I., Hofmeister, A. M., & Whittington, A. G. (2012). The influence of temperature-dependent thermal diffusivity on the conductive cooling rates of plutons and temperature-time paths in contact aureoles. *Earth and Planetary Science Letters*, 317–318, 157–164. <https://doi.org/10.1016/j.epsl.2011.11.009>

- Norton, D., & Knight, J. (1977). Transport phenomena in hydrothermal systems: Cooling plutons. *American Journal of Science*, 277(8), 937–981. <https://doi.org/10.2475/ajs.277.8.937>
- Norton, D., & Taylor, H. P. (1979). Quantitative simulation of the hydrothermal systems of crystallizing magmas on the basis of transport theory and oxygen isotope data: An analysis of the Skaergaard intrusion. *Journal of Petrology*, 20(3), 421–486. [https://doi.org/10.1093/](https://doi.org/10.1093/petrology/20.3.421)
- Parmentier, E. M., & Schedl, A. (1981). Thermal aureoles of igneous intrusions—Some possible indications of hydrothermal convective cooling. *The Journal of Geology*, 89(1), 1–22. <https://doi.org/10.1086/628561>
- Reiners, P. W., & Brandon, M. T. (2006). Using thermochronology to understand orogenic erosion. *Annual Review of Earth and Planetary Sciences*, 34(1), 419–466. <https://doi.org/10.1146/earth.2006.34.issue-1>
- Reiners, P. W., Ehlers, T. A., Mitchell, S. G., & Montgomery, D. R. (2003). Coupled spatial variations in precipitation and long-term erosion rates across the Washington cascades. *Nature*, 426(6967), 645–647. <https://doi.org/doi:10.1038/nature02111>
- Roy, M., Kelley, S., Pazzaglia, F., Cather, S., & House, M. (2004). Middle tertiary buoyancy modification and its relationship to rock exhumation, cooling, and subsequent extension at the eastern margin of the Colorado Plateau. *Geology*, 32(10), 925–928. [https://doi.org/10.1130/](https://doi.org/10.1130/G20561.1)
- Taylor, J. P., & Fitzgerald, P. G. (2011). Low-temperature thermal history and landscape development of the eastern Adirondack Mountains, New York: Constraints from apatite fission-track thermochronology and apatite (U-Th)/He dating. *Geological Society of America Bulletin*, 123(3–4), 412–426. <https://doi.org/10.1130/B30138.1>
- Thomson, S. N., Brandon, M. T., Tomkin, J. H., Reiners, P. W., Vásquez, C., & Wilson, N. J. (2010). Glaciation as a destructive and constructive control on mountain building. *Nature*, 467(7313), 313–317. <https://doi.org/10.1038/nature09365>
- Turcotte, D. L., & Schubert, G. (2002). *Geodynamics* (2nd ed.). New York: Cambridge University Press. [https://doi.org/10.1017/](https://doi.org/10.1017/CBO9780511807442)