



Originally published as:

Heinz, J., Waajen, A. C., Airo, A., Alibrandi, A., Schirmack, J., Schulze-Makuch, D. (2020): Bacterial Growth in Chloride and Perchlorate Brines: Halotolerances and Salt Stress Responses of *Planococcus halocryophilus*. - *Astrobiology*, 20, 2.

DOI: <http://doi.org/10.1089/ast.2019.2069>

Bacterial Growth in Chloride and Perchlorate Brines: Halotolerances and Salt Stress Responses of *Planococcus halocryophilus*

Jacob Heinz,¹ Annemiek C. Waajen,^{1,2} Alessandro Airo,¹ Armando Alibrandi,¹ Janosch Schirmack,¹ and Dirk Schulze-Makuch^{1,3,4,5}

Abstract

Extraterrestrial environments encompass physicochemical conditions and habitats that are unknown on Earth, such as perchlorate-rich brines that can be at least temporarily stable on the martian surface. To better understand the potential for life in these cold briny environments, we determined the maximum salt concentrations suitable for growth (MSCg) of six different chloride and perchlorate salts at 25°C and 4°C for the extremotolerant cold- and salt-adapted bacterial strain *Planococcus halocryophilus*. Growth was measured through colony-forming unit (CFU) counts, while cellular and colonial phenotypic stress responses were observed through visible light, fluorescence, and scanning electron microscopy. Our data show the following: (1) The tolerance to high salt concentrations can be increased through a stepwise inoculation toward higher concentrations. (2) Ion-specific factors are more relevant for the growth limitation of *P. halocryophilus* in saline solutions than single physicochemical parameters like ionic strength or water activity. (3) *P. halocryophilus* shows the highest microbial sodium perchlorate tolerance described so far. However, (4) MSCg values are higher for all chlorides compared to perchlorates. (5) The MSCg for calcium chloride was increased by lowering the temperature from 25°C to 4°C, while sodium- and magnesium-containing salts can be tolerated at 25°C to higher concentrations than at 4°C. (6) Depending on salt type and concentration, *P. halocryophilus* cells show distinct phenotypic stress responses such as novel types of colony morphology on agar plates and biofilm-like cell clustering, encrustation, and development of intercellular nanofilaments. This study, taken in context with previous work on the survival of extremophiles in Mars-like environments, suggests that high-concentrated perchlorate brines on Mars might not be habitable to any present organism on Earth, but extremophilic microorganisms might be able to evolve thriving in such environments. Key Words: Brines—Salt—Growth—Mars—Perchlorate—Halotolerance. Astrobiology 20, xxx–xxx.

1. Introduction

MOST SUBZERO SALINE HABITATS on Earth are dominated by sodium chloride (NaCl), and most research has been focused on this salt (for review, *e.g.*, Gunde-Cimerman *et al.*, 2018). However, non-NaCl saline environments exist on Earth as well, such as the calcium chloride-rich Don Juan Pond, Antarctica (Dickson *et al.*, 2013) or the Spotted Lake, Canada, having high sulfate concentrations (Pontefract *et al.*, 2017).

Similarly, soils on Mars contain non-NaCl salts such as perchlorates (Hecht *et al.*, 2009). Accordingly, the presence of perchlorate-rich martian groundwater has been discussed (Clifford *et al.*, 2010). Fitting well to this hypothesis, it was recently proposed that a discovered subsurface lake might contain magnesium and calcium perchlorates causing a freezing point depression of water down to its calculated temperature of –68°C (Orosei *et al.*, 2018). Furthermore, spectral investigations indicated that perchlorate salt hydrates and their brines might play a role

¹Center of Astronomy and Astrophysics, Astrobiology Research Group, Technical University of Berlin, Berlin, Germany.

²UK Centre for Astrobiology, School of Physics and Astronomy, University of Edinburgh, Edinburgh, UK.

³School of the Environment, Washington State University, Pullman, Washington, USA.

⁴GFZ German Center for Geoscience, Section Geomicrobiology, Potsdam, Germany.

⁵Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Department of Experimental Limnology, Stechlin, Germany.

in the formation of recurring slope lineae (RSL) on Mars (Ojha *et al.*, 2015).

At the Phoenix landing site, perchlorate concentrations in the martian soil are ranging from 0.4 to 0.6 wt% (Hecht *et al.*, 2009). However, it has to be kept in mind that the perchlorates are present as solid salts which will form highly concentrated brines whenever the temperature is above the eutectic temperature and the relative humidity (RH) is above the deliquescence relative humidity (DRH) of the perchlorate salt (Davila *et al.*, 2010; Nikolakakos and Whiteway, 2015; Heinz *et al.*, 2016). For example, at -34°C the sodium perchlorate concentration of the forming eutectic brine would be 52.6 wt% (9 M) (Hennings *et al.*, 2013), which is too high for any organism we know from Earth to thrive therein. Therefore, only diluted perchlorate brines might serve as a habitat. These diluted solutions could be stable in the subsurface of Mars (Burt, 2003; Martínez and Renno, 2013).

Since there are diurnal and seasonal temperature and humidity changes on Mars, it can be assumed that salt concentrations in these brines also fluctuate due to crystallization of ice or salt hydrates at cold temperature conditions or due to water absorption from the atmosphere at high RH conditions, for example in the morning hours when the RH in the martian atmosphere is highest and can reach saturation (Harri *et al.*, 2014). Thus, microorganisms would have to survive temporarily enhanced salt concentrations (including crystallization processes) at low temperatures and thrive at times of higher temperatures and brine dilution. It has already been shown that low temperatures enhance the bacterial survival in chloride (Cl^-) and perchlorate (ClO_4^-) brines with eutectic concentrations while, additionally, the high salt concentrations benefit the survival during freeze/thaw cycles of the brine (Heinz *et al.*, 2018). However, the question how low the salt concentration would have to be for microbial growth remained open prior to this study.

The exploration of the physicochemical limits for growth and survival of organisms thriving in cold saline environments gives not only insight into the microbial ecosystems adapted to the most extreme habitats on Earth but also provides the necessary data for assessing the habitability of extraterrestrial environments, for example on Mars (Schulze-Makuch *et al.*, 2015, 2017). A model organism for halo- and psychrotolerant bacteria is *Planococcus halocryophilus*, which has been isolated from the active layer of permafrost soil in the Canadian High Arctic (Myktyczuk *et al.*, 2012). It is able to grow at 19 wt/vol% NaCl (corresponding to 16 wt/wt%) concentration and at -15°C , while showing metabolic activity down to -25°C (Myktyczuk *et al.*, 2013; Raymond-Bouchard *et al.*, 2017).

Bacterial growth of *P. halocryophilus* under these harsh conditions is enabled by the expression of various osmolyte transporters and cold-adapted proteins, a high lipid turnover rate, and a high resource efficiency at subzero temperatures with an accumulation of carbohydrates as energy resource (Myktyczuk *et al.*, 2013). Furthermore, analyses of the proteome of *P. halocryophilus* revealed intricate changes in protein expression (Raymond-Bouchard *et al.*, 2017). Under subzero growth conditions, this bacterial strain develops a nodular sheet-like crust around the cells which might provide protection against cold and osmotic stress (Ronholm *et al.*, 2015; Myktyczuk *et al.*, 2016).

The ability to tolerate these cold and salty conditions was the reason for choosing *P. halocryophilus* as a suitable an-

alog microorganism to test how well microbes on Earth can adapt to the conditions prevailing on Mars and whether adaptation to high-concentration perchlorate brines is possible with the available biochemical toolset of life as we know it. *P. halocryophilus* is an aerobic organism and thus, at first glance, might not appear relevant to martian habitability, given that the molecular oxygen (O_2) content in the martian atmosphere is very low (0.13%). However, recent studies found that martian brines can be enriched with dissolved O_2 up to 2 mol m^{-3} , enabling aerobic microbes to metabolize (Stamenković *et al.*, 2018). In addition, there might be other suitable extraterrestrial habitats that provide both osmotic stress conditions and feasible O_2 levels.

Here, we investigated the maximum halotolerance for growth of *P. halocryophilus* at optimal growth temperature (25°C) and low temperature (4°C) for various Cl^- and ClO_4^- salts. Furthermore, we investigated the phenotypic adaptations to salt stress such as changes in cell and colony morphology and the formation of cell clusters. This study examines several major aspects important for astrobiological research especially on Mars where Cl^- and ClO_4^- brines might be the last possible refuges for life (Davila and Schulze-Makuch, 2016).

2. Materials and Methods

2.1. Organism and culture conditions

The bacterial strain *Planococcus halocryophilus* Or1 (DSM 24743^T), obtained from the DSMZ (Leibniz Institute DSMZ—German Collection of Microorganisms and Cell Cultures) was used in all experiments described within this study. The bacteria were grown aerobically at 25°C (optimum growth temperature) or 4°C (low temperature control) in liquid DSMZ growth medium #92 (3% Tryptic soy broth [TSB], 0.3% yeast extract) with various concentrations (1 wt% [w/w] incremental steps) of one of the following salts: sodium chloride (NaCl), magnesium chloride (MgCl_2), calcium chloride (CaCl_2), sodium perchlorate (NaClO_4), magnesium perchlorate ($\text{Mg}(\text{ClO}_4)_2$), or calcium perchlorate ($\text{Ca}(\text{ClO}_4)_2$). The media were prepared by mixing the media components, salt and water, followed by pH adjustment (pH 7.2–7.4), centrifugation if necessary (in calcium-containing samples, yeast flocculation can occur [Stratford, 1989] which has no influence on the cells' growth, because *P. halocryophilus* readily thrives in medium containing only TSB [Myktyczuk *et al.*, 2012]), and sterile filtration.

2.2. Determination of the maximum salt concentration suitable for growth (MSCg)

Bacteria were monitored for growth and death by using colony-forming unit (CFU) counts to determine the MSCg values of the respective salts. Two identical samples were prepared and inoculated separately (biological duplicates) for each experiment. For CFU counts, two aliquots of $100 \mu\text{L}$ were taken from each sample and plated on agar plates containing DSMZ growth medium #92. CFUs for the same experimental conditions were averaged by using the arithmetic mean. Where necessary, the aliquots were diluted with phosphate-buffered saline (PBS) containing additional 10 wt% NaCl (for all sodium-containing samples) or 10 wt% MgCl_2 (for all magnesium- or calcium-containing samples).

The increased amount of salt in the PBS was necessary to avoid bursting of cells by osmotic pressure during the dilution of the saline growth media. Some experiments were repeated to check reproducibility.

To investigate the effect of the inoculation culture on growth and survival of *P. halocryophilus* in the salty samples, 5 mL of the saline medium was inoculated with one of the following inoculation methods (IMs):

- IM 1: The medium was inoculated with the stock culture (growth medium + 10 wt% NaCl) at 25°C.
- IM 2: The medium was inoculated with a culture grown at the respective temperature (25°C or 4°C) in medium with lower concentration of the respective salt (progressive culture adaptation).
- IM 3: Medium for experiments at 25°C was inoculated with a culture grown at 4°C in medium with the same or higher concentration of the respective salt.
- IM 4: Medium for experiments at 4°C was inoculated with a culture grown at 25°C in medium with the same or higher concentration of the respective salt.

Inoculation volumes ranged from 10 to 50 μL depending on the cell density of the inoculation culture. However, due to cell clustering (Section 3.2.2) the starting cell density after inoculation varied between $5 \cdot 10^2$ CFU/mL and $5 \cdot 10^4$ CFU/mL.

Because progressive culture adaptations were done by a 1 wt% stepwise increase in the salt concentrations of the medium, the MSCg values had an inherent uncertainty of 1%. Larger uncertainties (up to 2%) occurred for some samples incubated at 4°C when cells in media with salt concentrations above the MSCg neither grew nor died within the time of the experiment.

2.3. Light, fluorescence, and scanning electron microscopy

A set of samples was investigated under the light microscope (Primo Star, Zeiss, equipped with Axio Cam 105 color) without prior sample treatment. For fluorescence microscopy of living and dead cells, samples were washed twice with PBS containing 10 wt% NaCl. Three milliliters of each sample was stained with 3 μL of a 1:2 mixture of component A (SYTO 9 dye, 3.34 mM) and component B (propidium iodide, 20 mM) from the Invitrogen Live/Dead BacLight Bacterial Viability Kit, where component A causes green fluorescence of intact cells and component B causes red or orange fluorescence of dead cells with damaged cell walls. The stained samples were imaged with a fluorescence microscope (Polyvar 2, Reichert-Jung) equipped with a Xenon lamp (XBO 150 W/1).

Samples for scanning electron microscopy (SEM) were washed twice with PBS containing 10 wt% NaCl followed by fixation in 2.5% glutaraldehyde solution (in 0.1 M phosphate buffer [PB], pH=7.3). The fixed samples were washed twice with 0.1 M PB, dehydrated through a graded acetone series (50%, 70%, 90%, 95%, 100%), critical point dried in a Leica CPD300, coated with carbon, and imaged with a Hitachi S-2700 microscope.

3. Results

3.1. Growth at 25°C and 4°C

3.1.1. Growth curves and MSCg values. For determining the MSCg values of all Cl^- and ClO_4^- salts, we used

growth versus death as a distinction criterion. For example, after 6 days of incubation in CaCl_2 -rich media, *P. halocryophilus* shows an increase in CFU values (*i.e.*, growth) at 25°C under all tested salt concentrations with the exception of 9 wt%; hence at this temperature the MSCg value is 8 wt% (Fig. 1). However, at 4°C the MSCg value is greater with 10 wt% (embedded plot of Fig. 1).

All growth curves generated are provided in the Supplementary Materials (Fig. S1–S12, available online at www.liebertonline.com/ast). The resulting MSCg values for all salts and temperatures, including their corresponding total ion concentrations (sum of cation and anion concentration), anion concentrations, ionic strengths, and water activities are listed in Table 1. The MSCg (wt%), total molar ion concentrations, and anion concentrations are plotted as bar charts in Fig. 2.

Overall, *P. halocryophilus* shows high halotolerances to all Cl^- and ClO_4^- salts at both 25°C and 4°C (Fig. 2A, 2B). The Cl^- tolerance is at least 2.5-fold higher than the tolerance to ClO_4^- in media with the same cation (Fig. 2C). However, with 12 wt% (1.1 M) NaClO_4 at 25°C we found the highest microbial tolerance to NaClO_4 described so far. The lowest tolerated water activity was 0.90 in 14 wt% (2.8 M) NaCl, while the highest tolerable ionic strength (3.9 mol/L) was reached in 11 wt% (1.3 M) MgCl_2 at 25°C (Table 1).

3.1.2. Effect of the inoculation method (IM) on the MSCg. The applied IM effects the growth curves and the resulting MSCg values in the following ways:

- (1) At 25°C, the MSCg values could only be reached with IM 2 (progressive culture adaptation at 25°C) but not with IM 1 (inoculation with stock culture) where cell death occurs already at lower salt concentrations. For example, the MSCg for MgCl_2 was 9 wt% at 25°C when the media was inoculated with the stock culture (IM 1); however, a stepwise (1 wt%) increase in the MgCl_2 concentration (IM 2) resulted in a MSCg of 11 wt% MgCl_2 (Fig. S3). It is notable that the length of the growth curve lag phase (occasionally including an initial CFU reduction) is enhanced with increasing salt concentration and decreasing temperatures (*e.g.*, Figs. 1 and S1).
- (2) Applying IM 3 (4°C \rightarrow 25°C inoculation) resulted in an increase of the MSCg values at 25°C only in the case of $\text{Ca}(\text{ClO}_4)_2$ samples. Here, growth in 3 wt% $\text{Ca}(\text{ClO}_4)_2$ was not detected after inoculation with the stock solution (IM 1) nor with a 2 wt% $\text{Ca}(\text{ClO}_4)_2$ culture grown at 25°C (IM 2), but only after inoculation with a 3 wt% $\text{Ca}(\text{ClO}_4)_2$ culture grown at 4°C (IM 3) (Fig. S11).
- (3) At 4°C, a higher MSCg value was reached by applying IM 2 (progressive culture adaptation at 4°C) than by applying IM 4 (25°C \rightarrow 4°C inoculation). For example, at 4°C inoculation of 2 wt% $\text{Mg}(\text{ClO}_4)_2$ medium with a 5 wt% $\text{Mg}(\text{ClO}_4)_2$ culture grown at 25°C did not show growth, indicating a MSCg < 2 wt% $\text{Mg}(\text{ClO}_4)_2$ when IM 4 is applied. However, a 1 wt% stepwise increase in $\text{Mg}(\text{ClO}_4)_2$ concentration at 4°C resulted in an culture able to growth at 4 wt% $\text{Mg}(\text{ClO}_4)_2$ (Fig. S10). These data suggest that for growth at 4°C an adaption to the cold first has to take place before *P. halocryophilus* can adapt stepwise to higher salt concentrations at that temperature.




FIG. 1. Growth curves of *P. halocryophilus* in liquid growth media with different CaCl_2 concentrations at 25°C (red curves) and 4°C (blue curves). X indicates the detection limit (no detectable CFUs in a 100 μL aliquot). IM describes the inoculation method as explained in Section 2.2. MSCg values are 8 and 10 wt% CaCl_2 for 25°C and 4°C, respectively. Growth in 10 wt% CaCl_2 at 4°C (embedded plot) was only observed for one of the two biological duplicates, thus, lacking an error bar.

3.1.3. Temperature effect on the MSCg. The relative shift in the MSCg that occurs by lowering the incubation temperature from 25°C to 4°C is visualized in Fig. 3. Among all six salts investigated in this study, only cells in CaCl_2 -containing media show an enhanced salt tolerance at lower temperature, where growth at 9 and 10 wt% CaCl_2 did not occur at 25°C but only at 4°C (Fig. 1).

Nevertheless, the observation that only IM 4 (4°C \rightarrow 25°C inoculation) resulted in growth at the MSCg for $\text{Ca}(\text{ClO}_4)_2$ (3 wt%) at 25°C (see point (2) of Section 3.1.2) provides evidence that also the tolerance to $\text{Ca}(\text{ClO}_4)_2$ is increased at 4°C, however, to a lower extent than the 1 wt% salt concentration incremental steps used in this study (Section 2.1). This suggests a general increase in the

TABLE 1. MSCG VALUES AND CORRESPONDING TOTAL ION CONCENTRATIONS (SUM OF CATIONS AND ANIONS), ANION CONCENTRATIONS, IONIC STRENGTHS, AND WATER ACTIVITIES AT 25°C AND 4°C

	MSCg		Total ion conc.		Anion conc.		Ionic strength		Water activity*			
	[wt%]		[mol/L]		[mol/L]		[mol/L]		[mol/L]			
	25°C	4°C	25°C	4°C	25°C	4°C	25°C	4°C	25°C	4°C		
NaCl	14 (1)	11 (1)	2.79	2.11	5.57	4.23	2.79	2.11	2.79	2.11	0.90	0.93
MgCl ₂	11 (1)	10 (2)	1.30	1.17	3.89	3.50	2.60	2.33	3.89	3.50	0.92	0.93
CaCl ₂	8 (1)	10 (1)	0.78	1.00	2.35	3.00	1.57	2.00	2.35	3.00	0.96	0.95
NaClO ₄	12 (1)	7 (2)	1.11	0.61	2.23	1.23	1.11	0.61	1.11	0.61	0.96	0.98
Mg(ClO ₄) ₂	5 (1)	4 (1)	0.24	0.19	0.71	0.56	0.47	0.37	0.71	0.56	0.99	0.99
Ca(ClO ₄) ₂	3 (1)	3 (1)	0.13	0.13	0.39	0.39	0.26	0.26	0.39	0.39	0.99	0.99

Values in parentheses give the deviation as described in Section 2.2.

*Water activity calculated from the Pitzer equation (Pitzer, 1991) with Pitzer parameters taken from Toner *et al.* (2015). The temperature dependence (25°C vs. 4°C) of the water activity is negligible for Cl^- (Fontan and Chirife, 1981) and ClO_4^- solutions (Toner and Catling, 2016) at temperatures above 0°C.




FIG. 2. Maximum salt concentrations suitable for growth (MSCg) of *P. halocryophilus* expressed as wt% (A), and total molar concentration (sum of cation and anion concentration) (B), and the molar anion concentrations at the corresponding MSCg (C). Transparent parts of the bars represent the salinity range for which neither growth nor complete demise of the culture could be determined within the time of the experiment (see Section 2.2). The water activities and ionic strengths at the MSCg are shown in Fig. S13.

calcium (Ca²⁺) tolerance of *P. halocryophilus* at lower temperatures.

In contrast, the sodium (Na⁺) tolerance is decreased at lower temperatures for both anions, Cl⁻ and ClO₄⁻ (Fig. 3). The tolerances to magnesium (Mg²⁺) are only slightly reduced at 4°C (1 wt% also for both anions, Cl⁻ and ClO₄⁻). The reduction in the Na⁺ tolerance at 4°C on the one hand and the increased Ca²⁺ tolerance at 4°C on the other hand led to an equalization of the anion (Cl⁻ or ClO₄⁻) concentration at

the MSCg at 4°C, while at 25°C the differences between the anion concentrations for the three different cationic species (Na⁺, Mg²⁺, and Ca²⁺) are more pronounced (Fig. 2).

3.2. Cellular and colonial phenotypic salt-stress adaptations

3.2.1. Colonial phenotypic adaptations. *P. halocryophilus* cells grown in liquid cultures exhibit with an increase in salt concentration, particularly for perchlorate salts, the tendency to develop macroscopic cohesive biofilms that could only be disrupted by intense shaking or vortexing (Fig. 4A).

Furthermore, it was observed that a novel colony phenotype (type II) appears only on plates inoculated with aliquots from cells grown in ClO₄⁻-rich medium (especially with >9 wt% NaClO₄ at 25°C). This type II colony is paler and duller than the usual colonies (type I) that are shiny and orange (Fig. 4) and does not occur in cultures grown in salt-free or Cl⁻-containing media. Occasionally, both colony types occurred on plates inoculated with aliquots from cells grown in media with perchlorate concentrations of a few weight percent below the MSCg (Fig. 4B). Sporadically, such colonies underwent a transformation from the type II back to the type I after approximately 2 weeks of growth on the agar plates (Fig. 4C). Contamination was ruled out through 16S sequencing of both colony types (99.90% sequence similarity of type I versus type II, data not shown), suggesting that the colony type II represents a reversible multigenerational phenotypic adaptation of *P. halocryophilus* to high perchlorate salt stress. Type II colonies needed 3–5 times longer than type I colonies to reach comparable colony sizes.

Two additional colony phenotypes were observed on agar plates: Type III colonies are irregular jagged in shape and occurred on agar plates inoculated with magnesium-rich (Cl⁻ and ClO₄⁻) cultures (Fig. S14). Type IV colonies are mucoid and shiny, merge rapidly during growth, and occurred on agar plates inoculated with CaCl₂-rich cultures (Fig. S15).

3.2.2. Cellular phenotypic adaptations. *P. halocryophilus* cells grown in liquid media containing no additional salts, seen under the light and fluorescence microscope as well as in SEM images, occur predominantly as single cells, diplococci, or small cell aggregates (Fig. 5A–5C) and have an overall smooth surface with nodules occurring largely along the cell division plane (Fig. 5B) as previously described (Mykytczuk *et al.*, 2016).

Cells, however, grown in Cl⁻-rich media occur predominantly in larger clusters of ~100 cells (Fig. 5D–5F); moreover, cells grown in perchlorate-rich media cluster into even larger aggregates of >1000 cells (Fig. 5G–5I). These clusters, containing living and dead cells (Fig. 5G), are highly cohesive and could not be disrupted by 5 min of ultrasonication nor by washing with 100% ethanol and killing all cells (Fig. S16). The cell clustering predominantly occurring in perchlorate-rich media led to higher uncertainties and irregularities in the growth curves of these samples (*e.g.*, Figs. S7 and S11). Concurrent with cell clustering under salt-stress conditions is the development of intercellular nanofilaments within a cluster (Fig. 5E). Additionally, cells in CaCl₂-rich media developed surface encrustation (Fig. 5F).




FIG. 3. Relative changes in the MSCg induced by lowering the incubation temperature from 25°C to 4°C.




FIG. 4. Macroscopically visible salt stress responses: (A) Biofilm-like cell clumping occurring in a 10 wt% NaClO₄ culture after 1 month of growth. (B) Two different colony morphologies of *P. halocryophilus* observed 1 week after plating a 9 wt% NaClO₄ culture at 25°C. The shiny intense-orange colonies (type I) represent the colony morphology typical for *P. halocryophilus* grown in medium with low salt concentrations. The paler and smaller colonies (type II) only occurred after plating perchlorate-rich cultures. (C) Transformation of a type II colony into type I after 2 weeks of colony growth.




FIG. 5. Light microscopy (A, D), fluorescence microscopy (after live/dead staining) (G), and SEM (B–C, E–F, H–I) images of *P. halocryophilus* after growth in media containing no salts (A–C), chlorides (D–F), and perchlorates (G–I) at 25°C. Cells grown in salt-free media developed smooth surfaces with some nodular features (n) and occurred as single cells, diplococci, or smaller cell aggregates (A–C). Several salt stress responses were observed including formation of cell clusters (G–I) and filaments (f) (E), and encrustation (en) of some cells in CaCl_2 -containing cultures (F).

4. Discussion

4.1. Salt-stress response and phenotypic adaptation

Microbial salt-stress responses and multigenerational adaptations such as microbial cell aggregation and biofilm formation have been observed for various stress conditions such as desiccation (Monier and Lindow, 2003), UV radiation (Fröls *et al.*, 2008), or NaCl exposure (Philips *et al.*, 2017). Thus far, NaCl is the only salt for which the salt-stress response of *P. halocryophilus* has been investigated, showing that the cells developed an EPS-like coating with short filament-like features if grown in 15 wt% (18 wt/vol%) NaCl media at -5°C (see Fig. 1e, 1f in Mykytczuk *et al.*, 2016).

For our experimental growth conditions, we show a similar trend for *P. halocryophilus* at high NaCl concentrations, where the cells developed even longer nanofilaments (Fig. 5E). The formation of similar nanofilaments linking cells within one cluster has also been observed previously, for example for the halophilic archaeon *Halo-coccus salifodinae* (see Fig. 2 in Legat *et al.*, 2013). Furthermore, we conducted experiments with two additional Cl^- and three ClO_4^- salts showing that *P. halocryophilus* develops particularly large and highly cohesive clusters

especially if grown in perchlorate-rich media (Fig. 5G–5I). Previous studies on *Hydrogenothermus marinus* have shown a formation of cell chains under increased perchlorate concentrations which has been speculated to result from incomplete cell division (Beblo-Vranesevic *et al.*, 2017), a physiological effect of perchlorate that could be similar for *P. halocryophilus*. Such a mechanism would be in accordance with our observation that cell clusters could neither be destroyed through ultrasonication nor through killing the cells with ethanol treatment (Fig. S16).

Furthermore, our results show the development of cohesive biofilms in perchlorate-rich media (Fig. 4A), and if transferred to agar plates, the occurrence of an additional colony morphology (type II, Fig. 4B). These are both macroscopic phenotypes consistent with the microscopic development of large cell clusters linked by numerous nanofilaments. Such stress-induced changes in colony morphology are also known for other stressors than high salinity; for example, under nutrient starvation *Vibrio cholerae* colonies change from the normal translucent to a rugose type (Wai *et al.*, 1998), or with a pH shift *Bacillus subtilis* colonies change from the normal notched “volcano-like” to a round and front-elevated “crater-like” shape (Tasaki *et al.*, 2017).

The phenotypic responses described above demonstrate that organisms like *P. halocryophilus* can develop perchlorate-specific stress adaptations that are not (or only to a lower extent) used to counteract high chloride concentrations. This is an important finding for all extraterrestrial environments with naturally occurring perchlorates. This might include not only Mars but any planetary bodies with a relatively dry surface (to avoid leaching of salts) and increased UV radiation (to oxidize chlorides [Carrier and Kounaves, 2015]). For example, spectral data indicate the presence of perchlorates at the surface of the icy moon Europa (Ligier *et al.*, 2016), which could entail delivery of perchlorates to Europa's subsurface ocean (Hand *et al.*, 2007). Also, based on the observed phenotypic adaptations of *P. halocryophilus*, microfossils of such organisms found in perchlorate-rich environments might be present in the form of cell clusters or biofilms rather than in the form of single cells.

4.2. Halotolerances of *P. halocryophilus* at 25°C and 4°C

To our knowledge, the only MSCg data for *P. halocryophilus* have been reported for NaCl being 16 wt% (19 wt/vol%) at 25°C (Mykytczuk *et al.*, 2012) and 15 wt% (18 wt/vol%) + 7 wt/vol% glycerol at -15°C (Mykytczuk *et al.*, 2013, 2016). In contrast, we found a lower NaCl halotolerance of 14 and 11 wt% at 25°C and 4°C, respectively, which can have various causes. For example: (1) Mutation of the lab culture and loss of part of its physiological abilities since isolation from its natural environment and first description (Mykytczuk *et al.*, 2012). (2) Different, and commonly not or only partially reported, pre-conditioning and adaptation procedures for cell growth such as the increment size of salt increase, growth curve phase used for transfer inoculation, and transfer or culture volume and agitation. (3) Use of a different growth media: while we used DSMZ growth media #92 containing TSB and yeast extract, Mykytczuk (2013 and 2016) used a growth media containing TSB and glycerol to maintain the medium liquid down to temperatures of -15°C. Glycerol is known to be an osmoprotectant and, thus, might have caused the elevated NaCl tolerance at low temperatures. Whether this difference in halotolerance would have also applied to other salts remains unknown, since no other salts than NaCl have been investigated for *P. halocryophilus* prior to this study.

The determination of MSCg values for multiple salts, as presented here for the first time for *P. halocryophilus*, provides the opportunity to test if any single physicochemical aspect of saline solutions is a limiting factor for growth. Our data, however, do not indicate that any individual factor (including total ion or anion concentration, ionic strength, water activity; see Table 1, Fig. 2, and Fig. S13) is responsible for the growth limitation of *P. halocryophilus* in saline solutions. Otherwise, the values of one of these physicochemical factors (*e.g.*, water activity) would be similar for all salts, which is not the case.

However, our data indicate the following ion species-dependent halotolerances for *P. halocryophilus*.

- (1) The anion species, here Cl⁻ versus ClO₄⁻, plays the most dominant role determining the MSCg values independent of physicochemical unit, showing an overall at least 2.5-fold higher tolerance to Cl⁻ than to ClO₄⁻ salts with the

same cation. Nevertheless, the MSCg of NaClO₄ (12 wt%; 1.1 M) is comparatively high and exceeds earlier findings for other organisms such as *Haloferax mediterranei* (0.6 M) (Oren *et al.*, 2014), *Halomonas venusta* (1.0 M) (Al Soudi *et al.*, 2017), *Hydrogenothermus marinus* (0.44 M) (Beblo-Vranesevic *et al.*, 2017), different bacterial isolates from Big Soda Lake in Nevada, USA (0.17 M) (Matsubara *et al.*, 2017), and *Halorubrum lacusprofundi* (0.8 M) (Laye and DasSarma, 2018). It should be noted that it is not clear how *P. halocryophilus* developed such high perchlorate tolerances. Perchlorates are rare in natural terrestrial environments and often coupled to the occurrences of nitrates, for example in the Atacama Desert, Chile (Dasgupta *et al.*, 2005). No nitrates have been detected in the permafrost samples *P. halocryophilus* has been isolated from (Steven *et al.*, 2007); thus it appears likely that this strain has never been exposed to environmental perchlorates and did not derive its resistance to perchlorate that way.

- (2) The cation species plays a similar role for growth at 25°C, where Na⁺ is the most tolerated cation for each group of anions, that is, NaCl among the chlorides and NaClO₄ among the perchlorates.

In summary, for growth at 25°C, Na⁺ and Cl⁻ ions are individually the most tolerated ions, and NaCl is the salt to which *P. halocryophilus* has the highest halotolerance (2.8 M) at the lowest water activity (0.90). Life on Earth has generally adapted most efficiently to the most abundant and/or common factors in nature; for salt, this is NaCl. Hence, our data suggest that the limitation of growth for *P. halocryophilus* is to a large degree based on evolutionary adaptations to brine veins (most likely consisting of NaCl) present in permafrost soil (Steven *et al.*, 2007). Also, all members of the genus *Planococcus* are halotolerant and have been isolated predominantly from cold and/or saline environments like the Arctic, Antarctic, or marine habitats (Mykytczuk *et al.*, 2012).

Perchlorate ions in aqueous solutions are relatively inert and non-oxidizing due to kinetic barriers (Urbansky, 1998) but rare in natural environments on Earth and therefore presumably exhibit an enhanced toxicity compared to chlorides. Thus, it seems plausible that putative martian microbes could adapt to naturally occurring perchlorate-rich environments to the same extent as Earth microbes such as *Planococcus* spp. did adapt to NaCl-enriched habitats. This idea is consistent with our finding that the tolerance to high salt concentrations can be increased through a stepwise inoculation toward higher concentrations. At low temperatures (4°C in this study) longer lag phases would provide even more time for adaption to higher salt concentrations to occur.

Additional to the ion species-dependent halotolerance, a temperature-dependent halotolerance of *P. halocryophilus* was observed, where at 25°C the MSCg values are different for each cation species, while at 4°C the cation species is less relevant and the MSCg values are more similar (Fig. 2). This MSCg value alignment at low temperatures appears to be largely caused by two separate trends.

- (1) Na⁺-containing salts (*i.e.*, NaCl and NaClO₄) can be tolerated by *P. halocryophilus* to higher concentrations at higher temperatures. Possibly, the elaborate biochemical machinery evolved to cope with high Na⁺ concentrations

is kinetically more effective at optimal growth temperatures enhancing the overall halotolerance.

- (2) In the case of CaCl_2 we observe the opposite effect, where lowering the temperature increases the halotolerance to this salt. This observation is in accordance with a previous study showing that with decreasing temperature the survival of *P. halocryophilus* in eutectic CaCl_2 brines is enhanced to a significantly larger degree than in NaCl , MgCl_2 , or NaClO_4 brines (Heinz *et al.*, 2018). A low temperature-induced halotolerance enhancement has been described for example for the bacterial strain *Clostridium* sp. isolated from brine lenses in the Siberian permafrost (Gilichinsky *et al.*, 2003) and for *M. soligelidi* (Morozova and Wagner, 2007) in NaCl -rich media. However, to our knowledge, *P. halocryophilus* is the first organism described thus far that shows an increased CaCl_2 tolerance at lower temperatures and can grow at salt concentrations up to 10 wt% (1 M) CaCl_2 at 4°C in the absence of kosmotropic ions which otherwise could compensate the chaotropic stress caused by calcium ions (Oren, 2013).

However, it remains unclear what mechanism causes the enhanced CaCl_2 tolerance of *P. halocryophilus* at 4°C, perhaps a psychrophilic optimization of the relevant biochemical machinery for coping with Ca^{2+} or simply the lethal effect of calcium being decreased at lower temperatures. It has been proposed that the increased CaCl_2 tolerance at lower temperatures might be due to the formation of larger and more stable hydration shells around calcium ions with decreasing temperature (Heinz *et al.*, 2018). A possible biological explanation is a beneficial effect caused by cellular encrustation, which was only observed in CaCl_2 -rich media in this study (Fig. 5F). Correspondingly, cellular encrustation containing 20% calcium carbonate has been documented previously for *P. halocryophilus* cells grown at subzero temperatures in NaCl -rich media (Mykytczuk *et al.*, 2016). Similar encrustation might be triggered in the presence of high Ca^{2+} amounts and might provide an efficient calcium resistance strategy due to the microbial mediated calcium carbonate precipitation. This positive effect might be increased at lower temperatures.

Another factor that might play a role for the enhanced Ca^{2+} tolerance at lower temperatures is the chaotropicity of Ca^{2+} . Chaotropic compounds increase the flexibility of macromolecules like proteins and, thus, can limit life at high temperatures (Hallsworth *et al.*, 2007), but they can also benefit microbial growth at low temperatures (Chin *et al.*, 2010; Rummel *et al.*, 2014). Cray *et al.* (2013) found that the chaotropic activity for CaCl_2 ($92.2 \text{ kJ kg}^{-1} \text{ mol}^{-1}$) is significantly higher than for MgCl_2 ($54.0 \text{ kJ kg}^{-1} \text{ mol}^{-1}$) while NaCl is a kosmotropic compound ($-11.0 \text{ kJ kg}^{-1} \text{ mol}^{-1}$). These data suggest that the high chaotropicity of CaCl_2 (and potentially also of $\text{Ca}(\text{ClO}_4)_2$, but data for perchlorates are lacking) might contribute to the enhanced microbial Ca^{2+} tolerance at low temperatures.

Additional work is needed to better understand the observed general trends in the habitability of Mars-analog brines in dependence of the type of salt, its concentration, and temperature. Additional long-term studies under subzero temperatures, similar to martian environments, should be conducted, especially since studies have shown that *P. halocryophilus* is able to grow under these conditions (Mykytczuk *et al.*, 2013,

2016). It is possible that lowering the experimental temperature to subzero values will further increase the tolerance of *P. halocryophilus* to Ca^{2+} . We also recommend that other microbial strains (including anaerobic ones) or communities should be investigated under similar experimental conditions to widen our understanding of life in cold brines.

5. Conclusion

For the first time, this study provides insights into the extremotolerant bacterium *P. halocryophilus*, which is well known for its tolerance to both cold temperatures and high concentration of salts, on how it survives and adapts not only to NaCl solutions but all Mars-relevant Cl^- and ClO_4^- salt solutions at different temperatures. Although growth in highly concentrated eutectic brines is not possible (Heinz *et al.*, 2018), the tolerance to the salts investigated in this study is intriguing and can even be enhanced by a stepwise increase in the salt concentration. For example, with 12 wt% (1.1 M) NaClO_4 we found the highest bacterial tolerance to ClO_4^- reported to date. For CaCl_2 -containing cell cultures we could show that by lowering the temperature from 25°C to 4°C the halotolerance increases from 8 wt% (0.8 M) to 10 wt% (1.0 M), respectively, while tolerances to Na^+ and Mg^{2+} are decreased for the same temperature decline.

The increased Ca^{2+} tolerance at lower temperatures corresponds well to the previously described enhanced survival in low-temperature brines (Heinz *et al.*, 2018) and plays an important role for the habitability of martian environments where Ca^{2+} -rich brines might be present in the shallow subsurface (Burt, 2003). Furthermore, Ca^{2+} is thought to be the main counter ion in ClO_4^- salts on Mars (Kounaves *et al.*, 2014), and $\text{Ca}(\text{ClO}_4)_2$ might be a main component of the recently discovered subsurface lake near the martian south pole and could be responsible for its freezing point depression (Orosei *et al.*, 2018).

Additionally, we described several salt adaption mechanisms like cell clustering, the formation of nanofilaments, encrustation of cells, and changes in the cell colony morphology. These data provide insight into how life could adapt to such high salt concentrations necessary for a sufficient freezing point depression allowing liquid water to be stable close to the martian surface.

Acknowledgments

This project was funded by European Research Council Advanced Grant “Habitability of Martian Environments” (HOME, no. 339231). All the numerical data for this paper are provided in the figures, tables, and supplementary information and are also available in tabular form from the authors upon request (schulze-makuch@tu-berlin.de). We would like express our thanks to two anonymous reviewers whose comments and suggestions helped improve this manuscript.

Author Disclosure Statement

No competing financial interests exist.

References

- Al Soudi, A.F., Farhat, O., Chen, F., Clark, B.C., and Schneegurt, M.A. (2017) Bacterial growth tolerance to concentrations of

- chlorate and perchlorate salts relevant to Mars. *Int J Astrobiol* 16: 229–235.
- Beblo-Vranesevic, K., Huber, H., and Rettberg, P. (2017) High tolerance of *Hydrogenothermus marinus* to sodium perchlorate. *Front Microbiol* 8, doi:10.3389/fmicb.2017.01369.
- Burt, D.M. (2003) Electrically conducting, Ca-rich brines, rather than water, expected in the martian subsurface. *J Geophys Res* 108, doi:10.1029/2002JE001862.
- Carrier, B.L. and Kounaves, S.P. (2015) The origins of perchlorate in the martian soil. *Geophys Res Lett* 42:3739–3745.
- Chin, J.P., Megaw, J., Magill, C.L., Nowotarski, K., Williams, J.P., Bhaganna, P., Linton, M., Patterson, M.F., Underwood, G.J.C., Mswaka, A.Y., and Hallsworth, J.E. (2010) Solutes determine the temperature windows for microbial survival and growth. *Proc Natl Acad Sci USA* 107:7835–7840.
- Clifford, S.M., Lasue, J., Heggy, E., Boisson, J., McGovern, P., and Max, M.D. (2010) Depth of the martian cryosphere: revised estimates and implications for the existence and detection of subpermafrost groundwater. *J Geophys Res* 115, doi:10.1029/2009JE003462.
- Cray, J.A., Russell, J.T., Timson, D.J., Singhal, R.S., and Hallsworth, J.E. (2013) A universal measure of chaotropy and kosmotropy. *Environ Microbiol* 15:287–296.
- Dasgupta, P.K., Martinelango, P.K., Jackson, W.A., Anderson, T.A., Tian, K., Tock, R.W., and Rajagopalan, S. (2005) The origin of naturally occurring perchlorate: the role of atmospheric processes. *Environ Sci Technol* 39:1569–1575.
- Davila, A.F. and Schulze-Makuch, D. (2016) The last possible outposts for life on Mars. *Astrobiology* 16:159–168.
- Davila, A.F., Dupont, L.G., Melchiorri, R., Jänchen, J., Valea, S., de los Rios, A., Fairén, A.G., Möhlmann, D., McKay, C.P., Ascaso, C., and Wierzbos, J. (2010) Hygroscopic salts and the potential for life on Mars. *Astrobiology* 10:617–628.
- Dickson, J.L., Head, J.W., Levy, J.S., and Marchant, D.R. (2013) Don Juan Pond, Antarctica: near-surface CaCl₂-brine feeding Earth's most saline lake and implications for Mars. *Sci Rep* 3, doi:10.1038/srep01166.
- Fontan, C.F. and Chirife, J. (1981) The evaluation of water activity in aqueous solutions from freezing point depression. *Int J Food Sci Technol* 16:21–30.
- Fröls, S., Ajon, M., Wagner, M., Teichmann, D., Zolghadr, B., Folea, M., Boekema, E.J., Driessen, A.J.M., Schleper, C., and Albers, S.-V. (2008) UV-inducible cellular aggregation of the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by pili formation. *Mol Microbiol* 70:938–952.
- Gilichinsky, D., Rivkina, E., Shcherbakova, V., Laurinavichuis, K., and Tiedje, J. (2003) Supercooled water brines within permafrost—an unknown ecological niche for microorganisms: a model for astrobiology. *Astrobiology* 3:331–341.
- Gunde-Cimerman, N., Plemenitaš, A., and Oren, A. (2018) Strategies of adaptation of microorganisms of the three domains of life to high salt concentrations. *FEMS Microbiol Rev* 42:353–375.
- Hallsworth, J.E., Yakimov, M.M., Golyshin, P.N., Gillion, J.L.M., D'Auria, G., de Lima Alves, F., La Cono, V., Genovese, M., McKew, B.A., Hayes, S.L., Harris, G., Giuliano, L., Timmis, K.N., and McGenity, T.J. (2007) Limits of life in MgCl₂-containing environments: chaotropy defines the window. *Environ Microbiol* 9:801–813.
- Hand, K.P., Carlson, R.W., and Chyba, C.F. (2007) Energy, chemical disequilibrium, and geological constraints on Europa. *Astrobiology* 7:1006–1022.
- Harri, A.-M., Genzer, M., Kempainen, O., Gomez-Elvira, J., Haberle, R., Polkko, J., Savijärvi, H., Rennó, N., Rodriguez-Manfredi, J.A., Schmidt, W., Richardson, M., Siili, T., Paton, M., La Torre-Juarez, M.D., Mäkinen, T., Newman, C., Rafkin, S., Mischna, M., Merikallio, S., Haukka, H., Martin-Torres, J., Komu, M., Zorzano, M.-P., Peinado, V., Vazquez, L., and Urqui, R. (2014) Mars Science Laboratory relative humidity observations: initial results. *J Geophys Res Planets* 119:2132–2147.
- Hecht, M.H., Kounaves, S.P., Quinn, R.C., West, S.J., Young, S.M.M., Ming, D.W., Catling, D.C., Clark, B.C., Boynton, W.V., Hoffman, J., Deflores, L.P., Gospodinova, K., Kapit, J., and Smith, P.H. (2009) Detection of perchlorate and the soluble chemistry of martian soil at the Phoenix lander site. *Science* 325:64–67.
- Heinz, J., Schulze-Makuch, D., and Kounaves, S.P. (2016) Deliquescence-induced wetting and RSL-like darkening of a Mars analogue soil containing various perchlorate and chloride salts. *Geophys Res Lett* 43:4880–4884.
- Heinz, J., Schirmack, J., Airo, A., Kounaves, S.P., and Schulze-Makuch, D. (2018) Enhanced microbial survivability in subzero brines. *Astrobiology* 18:1171–1180.
- Hennings, E., Heinz, J., Schmidt, H., and Voigt, W. (2013) Freezing and hydrate formation in aqueous sodium perchlorate solutions. *Z Anorg Allg Chem* 639:922–927.
- Kounaves, S.P., Chaniotakis, N.A., Chevrier, V.F., Carrier, B.L., Folds, K.E., Hansen, V.M., McElhoney, K.M., O'Neil, G.D., and Weber, A.W. (2014) Identification of the perchlorate parent salts at the Phoenix Mars landing site and possible implications. *Icarus* 232:226–231.
- Laye, V.J. and DasSarma, S. (2018) An Antarctic extreme halophile and its polyextremophilic enzyme: effects of perchlorate salts. *Astrobiology* 18:412–418.
- Legat, A., Denner, E.B.M., Dornmayr-Pfaffenhuemer, M., Pfeiffer, P., Knopf, B., Claus, H., Gruber, C., König, H., Wanner, G., and Stan-Lotter, H. (2013) Properties of *Halococcus salifodinae*, an isolate from Permian rock salt deposits, compared with halococci from surface waters. *Life* 3: 244–259.
- Ligier, N., Poulet, F., Carter, J., Brunetto, R., and Gourgéot, F. (2016) Vlt/sinfoni observations of Europa: new insights into the surface composition. *Astron J* 151, doi:10.3847/0004-6256/151/6/163.
- Martínez, G.M. and Renno, N.O. (2013) Water and brines on Mars: current evidence and implications for MSL. *Space Sci Rev* 175:29–51.
- Matsubara, T., Fujishima, K., Saltikov, C.W., Nakamura, S., and Rothschild, L.J. (2017) Earth analogues for past and future life on Mars: isolation of perchlorate resistant halophiles from Big Soda Lake. *Int J Astrobiol* 16:218–228.
- Monier, J.-M. and Lindow, S.E. (2003) Differential survival of solitary and aggregated bacterial cells promotes aggregate formation on leaf surfaces. *Proc Natl Acad Sci USA* 100: 15977–15982.
- Morozova, D. and Wagner, D. (2007) Stress response of methanogenic archaea from Siberian permafrost compared with methanogens from nonpermafrost habitats. *FEMS Microbiol Ecol* 61:16–25.
- Mykytczuk, N.C.S., Wilhelm, R.C., and Whyte, L.G. (2012) *Planococcus halocryophilus* sp. nov., an extreme sub-zero species from high Arctic permafrost. *Int J Syst Evol Microbiol* 62:1937–1944.
- Mykytczuk, N.C.S., Foote, S.J., Omelon, C.R., Southam, G., Greer, C.W., and Whyte, L.G. (2013) Bacterial growth at –15°C; molecular insights from the permafrost bacterium *Planococcus halocryophilus* Or1. *ISME J* 7:1211–1226.

- Mykytczuk, N.C.S., Lawrence, J.R., Omelon, C.R., Southam, G., and Whyte, L.G. (2016) Microscopic characterization of the bacterial cell envelope of *Planococcus halocryophilus* Or1 during subzero growth at -15°C . *Polar Biol* 39:701–712.
- Nikolakakos, G. and Whiteway, J.A. (2015) Laboratory investigation of perchlorate deliquescence at the surface of Mars with a Raman scattering lidar. *Geophys Res Lett* 42:7899–7906.
- Ojha, L., Wilhelm, M.B., Murchie, S.L., McEwen, A.S., Wray, J.J., Hanley, J., Massé, M., and Chojnacki, M. (2015) Spectral evidence for hydrated salts in recurring slope lineae on Mars. *Nat Geosci* 8:829–832.
- Oren, A. (2013) Life in magnesium- and calcium-rich hypersaline environments: salt stress by chaotropic ions. In *Poly-extremophiles: Life under Multiple Forms of Stress. Cellular Origin, Life in Extreme Habitats and Astrobiology*, edited by J. Seckbach, A. Oren, and H. Stan-Lotter, Springer Science and Business Media, Dordrecht, the Netherlands, pp 217–232.
- Oren, A., Eleri Bardavid, R., and Mana, L. (2014) Perchlorate and halophilic prokaryotes: implications for possible halophilic life on Mars. *Extremophiles* 18:75–80.
- Orosei, R., Lauro, S.E., Pettinelli, E., Cicchetti, A., Coradini, M., Cosciotti, B., Di Paolo, F., Flamini, E., Mattei, E., Pajola, M., Soldovieri, F., Cartacci, M., Cassenti, F., Frigeri, A., Giuppi, S., Martufi, R., Masdea, A., Mitri, G., Nenna, C., Noschese, R., Restano, M., and Seu, R. (2018) Radar evidence of subglacial liquid water on Mars. *Science* 361:490–493.
- Philips, J., Rabaey, K., Lovley, D.R., and Vargas, M. (2017) Biofilm formation by *Clostridium ljungdahlii* is induced by sodium chloride stress: experimental evaluation and transcriptome analysis. *PLoS One* 12, doi:10.1371/journal.pone.0170406.
- Pitzer, K.S. (1991) *Activity Coefficients in Electrolyte Solutions*, CRC Press, Boca Raton, FL.
- Pontefract, A., Zhu, T.F., Walker, V.K., Hepburn, H., Lui, C., Zuber, M.T., Ruvkun, G., and Carr, C.E. (2017) Microbial diversity in a hypersaline sulfate lake: a terrestrial analog of ancient Mars. *Front Microbiol* 8, doi:10.3389/fmicb.2017.01819.
- Raymond-Bouchard, I., Chourey, K., Altshuler, I., Iyer, R., Hettich, R.L., and Whyte, L.G. (2017) Mechanisms of subzero growth in the cryophile *Planococcus halocryophilus* determined through proteomic analysis. *Environ Microbiol* 19:4460–4479.
- Ronholm, J., Raymond-Bouchard, I., Creskey, M., Cyr, T., Cloutis, E.A., and Whyte, L.G. (2015) Characterizing the surface-exposed proteome of *Planococcus halocryophilus* during cryophilic growth. *Extremophiles* 19:619–629.
- Rummel, J.D., Beaty, D.W., Jones, M.A., Bakermans, C., Barlow, N.G., Boston, P.J., Chevrier, V.F., Clark, B.C., de Vera, J.-P.P., Gough, R.V., Hallsworth, J.E., Head, J.W., Hipkin, V.J., Kieft, T.L., McEwen, A.S., Mellon, M.T., Mikucki, J.A., Nicholson, W.L., Omelon, C.R., Peterson, R., Roden, E.E., Sherwood Lollar, B., Tanaka, K.L., Viola, D., and Wray, J.J. (2014) A new analysis of Mars “Special Regions”: findings of the second MEPAG Special Regions Science Analysis Group (SR-SAG2). *Astrobiology* 14:887–968.
- Schulze-Makuch, D., Schulze-Makuch, A., and Houtkooper, J.M. (2015) The physical, chemical and physiological limits of life. *Life* 5:1472–1486.
- Schulze-Makuch, D., Airo, A., and Schirmack, J. (2017) The adaptability of life on Earth and the diversity of planetary habitats. *Front Microbiol* 8, doi:10.3389/fmicb.2017.02011.
- Stamenković, V., Ward, L.M., Mischna, M., and Fischer, W.W. (2018) O_2 solubility in martian near-surface environments and implications for aerobic life. *Nat Geosci* 11:905–909.
- Steven, B., Briggs, G., McKay, C.P., Pollard, W.H., Greer, C.W., and Whyte, L.G. (2007) Characterization of the microbial diversity in a permafrost sample from the Canadian high Arctic using culture-dependent and culture-independent methods. *FEMS Microbiol Ecol* 59:513–523.
- Stratford, M. (1989) Yeast flocculation: calcium specificity. *Yeast* 5:487–496.
- Tasaki, S., Nakayama, M., and Shoji, W. (2017) Self-organization of bacterial communities against environmental pH variation: controlled chemotactic motility arranges cell population structures in biofilms. *PLoS One* 12, doi:10.1371/journal.pone.0173195.
- Toner, J.D. and Catling, D.C. (2016) Water activities of NaClO_4 , $\text{Ca}(\text{ClO}_4)_2$ and $\text{Mg}(\text{ClO}_4)_2$ brines from experimental heat capacities: water activity >0.6 below 200 K. *Geochim Cosmochim Acta* 181:164–174.
- Toner, J.D., Catling, D.C., and Light, B. (2015) Modeling salt precipitation from brines on Mars: evaporation versus freezing origin for soil salts. *Icarus* 250:451–461.
- Urbansky, E.T. (1998) Perchlorate chemistry: implications for analysis and remediation. *Bioremediat J* 2:81–95.
- Wai, S.N., Mizunoe, Y., Takade, A., Kawabata, S.I., and Yoshida, S.I. (1998) *Vibrio cholerae* O1 strain TSI-4 produces the exopolysaccharide materials that determine colony morphology, stress resistance, and biofilm formation. *Appl Environ Microbiol* 64:3648–3655.

Address correspondence to:

Dirk Schulze-Makuch
Center of Astronomy and Astrophysics
Astrobiology Research Group
Technical University of Berlin
Hardenbergstr. 36
10623 Berlin
Germany

E-mail: schulze-makuch@tu-berlin.de

Submitted 15 March 2019

Accepted 26 June 2019

Abbreviations Used

- CFU = colony-forming unit
DRH = deliquescence relative humidity
DSMZ = German Collection of Microorganisms and Cell Cultures
EPS = extracellular polymeric substance
IM = inoculation method
MSCg = maximum salt concentration suitable for growth
PB = phosphate buffer
PBS = phosphate-buffered saline
RH = relative humidity
SEM = scanning electron microscopy
TSB = Tryptic soy broth