

Hawkings, J. (2021): Trickle and Treat? The Critical Role of Marine-Terminating Glaciers as Icy Macronutrient Pumps in Polar Regions. - Journal of Geophysical Research: Biogeosciences, 126, 10, e2021JG006598.

<https://doi.org/10.1029/2021JG006598>

JGR Biogeosciences

COMMENTARY

10.1029/2021JG006598

Key Points:

- A new article by Williams et al. published in *JGR: Biogeosciences* prompts discussion of the role glaciers play in coastal nutrient cycling
- Marine-terminating glaciers play an important role in coastal ecosystems by indirectly supplying macronutrients to surface waters
- Substantial changes to marine nutrient cycling expected in glaciated regions as ice cover wanes in response to atmospheric and oceanic warming

Correspondence to:

J. R. Hawkings,
hawkings@sas.upenn.edu

Citation:

Hawkings, J. R. (2021). Trickle and treat? The critical role of marine-terminating glaciers as icy macronutrient pumps in polar regions. *Journal of Geophysical Research: Biogeosciences*, 126, e2021JG006598. <https://doi.org/10.1029/2021JG006598>

Received 25 AUG 2021

Accepted 28 SEP 2021

Trickle and Treat? The Critical Role of Marine-Terminating Glaciers as Icy Macronutrient Pumps in Polar Regions

Jon R. Hawkings^{1,2} 

¹Department of Earth and Environmental Science, University of Pennsylvania, Philadelphia, PA, USA, ²Interface Chemistry, GFZ-Potsdam, Potsdam, Germany

Abstract Nutrient supply to the coastal euphotic zone is critical in sustaining the productivity of ecosystems and provides an important feedback on the carbon cycle via stimulating the biological pump. The supply of coastal nutrients is typically thought to be dominated by terrestrial inputs and upwelling of deep ocean waters. Marine-terminating glaciers can provide nutrients via both mechanisms but the relative importance of each is dependent on complex physical–biogeochemical processes ranging from those in the glacier drainage system to the ice–ocean interface. Constraining these glacial nutrient supply processes in a remote part of the Canadian Arctic Archipelago (CAA) is the focus of a new study by Williams et al. (2021, <https://doi.org/10.1029/2021JG006289>). Their findings advance our understanding of the role glaciers play in polar nutrient cycling by highlighting significant shallow meltwater pumping of macronutrient-replete marine waters by CAA marine-terminating glaciers into the euphotic zone.

Plain Language Summary Glaciers that flow into the ocean supply essential nutrients for life to coastal ecosystems. The glacial meltwater itself provides a direct “trickle” supply. Pumping of macronutrient-rich ocean waters by upwelling of fresh meltwater when it exits the glacier provides an indirect nutrient “treat.” These glacially derived nutrients likely help maintain ecosystem health, but the mobilization and delivery processes involved are complex and poorly resolved, in part due to a lack of spatial and temporal data. Recent research, some of which is featured in the current issue of *Journal of Geophysical Research: Biogeosciences*, has helped to highlight the importance of marine-terminating glaciers in indirect nutrient supply by detailing the importance of the meltwater pump. Melting of glaciers under climate warming scenarios is likely to change the balance of and processes contributing to nutrient supply and therefore the health of societally important polar ecosystems.

The delivery of the macronutrients carbon (C), nitrogen (N), phosphorus (P), and silicon (Si) to the euphotic zone of the coastal ocean represents one of the most important biogeochemical processes on Earth (Jickells, 1998; Moore et al., 2013). Macronutrients sustain microbial primary and secondary production, which are vital components of the global carbon cycle (De La Rocha & Passow, 2014). These microscopic organisms form the base of the marine food web and therefore dictate the structure of the marine ecosystem, including the vitality of environmentally and economically important fisheries (Chassot et al., 2010). Both land inputs and the hydrodynamics of coastal waters play critical roles in replenishing surface waters with nutrients for growth (Tseng et al., 2014). The main sources of nutrients transported from land to ocean are rivers (Statham, 2012), groundwater (Santos et al., 2021), and atmospheric aerosols/dust (Jickells et al., 2016). Upwelling waters at continental and island margins bring nutrient-rich deep water to the ocean surface (Bruland et al., 2001; Gove et al., 2016; Watanabe et al., 2017). More recently, glaciers have been identified as potential macronutrient sources to coastal waters (J. Hawkings et al., 2016; Hendry et al., 2019; Vick-Majors et al., 2020). Far from the inhospitable and undynamic systems they are often perceived, glaciers host diverse microbial communities (Anesio et al., 2017) and sustain high rates of biogeochemical activity through the unique combination of liquid water and access to an abundance of finely comminuted and amorphous particles (glacial flour) generated from glacier sliding (Tranter & Wadham, 2014; Wadham et al., 2019). Recently observed and predicted rapid ice mass loss due to climatic warming (Hugonnet et al., 2021; Mougnot et al., 2019) highlights the need to better constrain the role of glaciers as dynamic components of coastal nutrient cycling.




Figure 1. A schematic of the current hypothesized physiochemical processes occurring at the ice–ocean aquatic critical zone. Two main nutrient delivery pathways are identified: (i) direct input of dissolved macronutrients from subglacial meltwater and (ii) indirect macronutrient-replete deep ocean water entrainment in the upwelling meltwater plume. The size of the arrows indicates approximate dissolved macronutrient flux terms. Other processes are also likely to imprint on the main two delivery pathways, but these remain poorly constrained. Figure adapted from Chu (2014).

The coastal regions proximal to significant ice cover harbor economically important ecosystems (Long & Jones, 2021). Circumstantial evidence elucidating the importance of glacially derived nutrients for these ecosystems has existed for over 50 years (Apolloni, 1973; Burton & Liss, 1973), including early observations of enhanced biological activity in marine waters proximal to glacier meltwater outflow dating to the 1930s (Hartley & Dunbar, 1938), and more recent studies identifying marine-terminating glacier fronts as foraging hotspots and refugia for marine life (Laidre et al., 2016; Lydersen et al., 2014; Nishizawa et al., 2019; Urban-ski et al., 2017). Glaciers deliver macronutrients to the euphotic zone of coastal waters via two broad mechanisms (Figure 1). First, the direct input of dissolved and labile particulate nutrients carried by meltwaters (J. R. Hawkings et al., 2015; Hood & Scott, 2008; J. B. Martin et al., 2020) and icebergs (Smith et al., 2013; Wadham et al., 2019). Second, indirect nutrient-rich deep ocean water entrainment in buoyant fresh meltwater plumes emerging from the subglacial hydrological system of marine-terminating glaciers (also commonly referred to as tidewater glaciers) and melting keel of icebergs (“meltwater pumping”; e.g., Cape, Straneo, et al., 2019; Cape, Vernet, et al., 2019; Greisman, 1979; Halbach et al., 2019; Hopwood et al., 2018; Kanna et al., 2018; Meire et al., 2017; Moon et al., 2018). Grounding line depth, that is, depth in the water column at which the glacier separates from the bedrock beneath it and where the majority of meltwater exits the glacier, is argued to be critical in determining the effectiveness of this meltwater pump (Hopwood et al., 2018). The impacts of these two delivery mechanisms are spatially and temporally complex and a topic of healthy debate in the literature.

There are very few studies linking land and marine measurements in an ice-to-ocean gradient, which has made accurately quantifying direct versus indirect macronutrient inputs from glaciers difficult. In this issue of *Journal of Geophysical Research: Biogeosciences*, Williams et al. (2021) present macronutrient

concentration data over a gradient from the marine-terminating Sverdrup Glacier to coastal waters up to ~25 km from the glacier terminus. The study highlights the impact of direct meltwater macronutrient export versus indirect meltwater entrainment of marine waters. The authors conducted their work in the heavily glaciated Canadian Arctic Archipelago (CAA) where no data have been collected since the early 1970s (Apolloni, 1973), apart from that in a companion article published concurrently in *Global Biogeochemical Cycles* (Bhatia et al., 2021). The study is one of the first to use an ice-to-ocean biogeochemical data set to provide evidence of significant nutrient entrainment from a shallow (approximately tens of meters depth) marine-terminating glacier (e.g., Figure 1). These data are highly valuable in constraining flows and fluxes of macronutrients to coastal ecosystems in glaciated regions and can be used to infer how these interface zones are likely to respond to future glacier retreat and melt in a warming climate.

The ice-to-ocean biogeochemical gradient sampled by Williams et al. (2021) provides several important insights into the role of marine-terminating glaciers in regional macronutrient cycling. The authors found dissolved nutrient concentrations (inorganic N as $\text{NO}_3^- + \text{NH}_4^+$, soluble reactive phosphorus [SRP], dissolved silicon, and dissolved organic carbon [DOC]) indicative of oligotrophic conditions in glacial meltwaters sampled at ice margin and surface locations (supraglacial ice, snow, and fresh surface melt). Nutrient concentrations suggested a depleted glacial meltwater endmember consistent with inputs predominantly from atmospheric deposition onto the ice surface (e.g., J. Hawkings et al., 2016; McCutcheon et al., 2021; Wadham et al., 2016). These low dissolved nutrient concentrations are inferred to be representative of the bulk meltwater discharge and impart a dilution effect upon mixing with marine waters at the ice–ocean interface.

The authors highlight a potential role for glacial DOC, in contrast to inorganic N, P, and Si. The glacially derived dissolved organic matter (DOM) exhibits fluorophores characteristic of microbially produced highly labile compounds (at least ~20% more protein-like fluorescence than marine samples; Dubnick et al., 2020), despite low overall concentrations of DOC (mostly <20 μM), compared to more recalcitrant marine organic matter dominated by humic-like fluorescence. Bioavailable glacial DOM is consistent with findings on Arctic land terminating glacier systems (Fellman et al., 2010; Kellerman et al., 2020; Pain et al., 2020), but this is the first demonstration of labile DOM delivery from a marine-terminating glacier. Distinctive DOM fluorescence in glacially modified coastal waters around southwestern Greenland has also been reported recently and may be linked to glacier DOM delivery across boundary currents, although the ultimate source is unclear (Hendry et al., 2021). It is interesting to speculate how large the impact of glacial DOM might be versus other local sources (e.g., primary production, upwelling, and nonglacial terrestrial inputs) and how rapidly each may be consumed by heterotrophic organisms in the marine environment, given the low export flux may be balanced by high bioavailability.

The nutrient concentrations in the study by Williams et al. (2021) are generally lower (up to more than an order of magnitude for lithogenic nutrients) than those reported in glacial meltwaters elsewhere. These low nutrient concentrations indicate limited interaction of meltwater with sediments beneath the glacier, highlighting the importance of the subglacial hydrological regime in nutrient mobilization (Irvine-Fynn et al., 2011). Sverdrup Glacier is a slow moving polythermal glacier, likely with extensive regions of cold-based ice (Wyatt & Sharp, 2015). In cold-based regions, the glacier is frozen to the bedrock and production of glacial flour is significantly lower than glaciers with extensive regions of fast flowing warm-based ice (e.g., southwest Greenland; Nienow et al., 2017). Sverdrup Glacier likely lacks the widespread subglacial drainage networks that promote water–rock interaction, biogeochemical weathering, and higher nutrient concentrations, particularly rock-derived nutrients like Si (e.g., J. R. Hawkings et al., 2017; Meire et al., 2016, where dissolved Si concentrations can exceed 50 μM in glacial meltwaters vs. ~detection in this study). Even with these differences, glacial meltwater in all systems studied appears mostly to provide a “trickle” of direct dissolved macronutrients to coastal ecosystems that can only sustain a small portion of the observed primary productivity.

The real “treat” from marine-terminating glaciers comes indirectly from meltwater-driven upwelling of nutrient-rich marine waters. Williams et al. (2021) corroborate a range of studies from the Arctic where likely >90% of NO_3^- and SRP in the euphotic zone is sourced from indirect meltwater pumping (Cape, Straneo, et al., 2019; Halbach et al., 2019; Kanna et al., 2018; Meire et al., 2017). Large marine-terminating glaciers in Greenland are estimated to provide a flux of dissolved macronutrients to the coastal euphotic

zone that rivals the major Arctic river inputs in magnitude (Cape, Straneo, et al., 2019), while even smaller marine-terminating glaciers provide a significant nutrient addition to coastal surface waters (e.g., Bowdoin Glacier, Kanna et al., 2018). These large glacier dissolved macronutrient subsidies are linked to enhanced proximal summertime productivity, and likely impact higher trophic levels (e.g., higher observed catch in coastal systems with significant marine-terminating glacier inputs; Meire et al., 2017) with clear relevance for ecosystem services.

The impact of meltwater-induced oceanic water entrainment on dissolved macronutrient availability (nitrogen and phosphorus) has been modeled by Hopwood et al. (2018) and is expected to be approximately 2 orders of magnitude greater than the “trickle” from direct meltwater input. The same study hypothesized that retreat of marine-terminating glaciers into shallower waters would suppress nutrient entrainment (the “productivity continuum”), as pumping theoretically occurs above the hypothesized nutricline. The anticipated net effect is that shallow meltwater pumping has a minimal impact on euphotic zone primary productivity.

Marine-terminating glaciers in the CAA have shallow grounding lines with subglacial meltwater discharging at depths of approximately tens meters. The study of Williams et al. (2021) therefore provides an ideal testing ground for the hypothesis that shallow meltwater pumping (~30 m) of intermediate oceanic waters is relatively unimportant for marine ecosystem productivity. Williams et al. (2021) found that meltwater pumping at Sverdrup Glacier still drives significant nutrient entrainment, similar to the findings from Halbach et al. (2019) in Kongsfjorden, Svalbard, with turbid plumes positively correlated to higher chlorophyll *a* concentration. The nutrient enrichment is more modest than deeply ground marine-terminating glaciers (e.g., ~5 μM of NO_3^- compared to ~10 μM in Nuup Kangerlua, Sermilik Fjord, and Bowdoin Fjord; Cape, Straneo et al., 2019; Kanna et al., 2018; Meire et al., 2017) but is somewhat surprising considering the expected productivity decline with decreasing grounding line depth. Indeed, shallow grounded marine-terminating glaciers in the remote CAA have the capacity to pump oceanic nutrient-rich water to the surface at a scale that rivals direct nutrient inputs from the largest Arctic river in North America (Mackenzie River; Williams et al., 2021). Glaciers in the CAA can therefore maintain a significant nutrient pump to surface waters at the ice–ocean interface, with importance for regional coastal ecosystem health.

Despite the progress in understanding the ice–ocean biogeochemical critical zone, there are still multiple uncertainties that exist. Even though the direct input of dissolved macronutrients from glaciers into the marine environment appears minor, glaciers may play an important role in micronutrient (trace elements essential for life) input. Glacial meltwaters in both polar regions appear enriched in particulate and filterable micronutrients (e.g., Fe, Mn, and Co), particularly where an extensive subglacial drainage system exists (Aciego et al., 2015; Bhatia et al., 2013; J. R. Hawkings et al., 2020). Evidence also suggests that glacially derived micronutrients are exported into the euphotic zone of coastal waters, where they are important in maintaining elevated concentrations, particularly of Fe (e.g., Arrigo et al., 2017; Bhatia et al., 2021; Cape, Straneo, et al., 2019; Herraiz-Borreguero et al., 2016; Hopwood et al., 2016; Kanna et al., 2020; Krisch et al., 2021). This glacier micronutrient “treat” is likely have disproportionate biogeochemical effects in the Arctic and Antarctic due to the differences in marine nutrient limitation patterns (Hopwood et al., 2020; Moore et al., 2013). For example, Southern Ocean waters are mostly Fe (and possibly Mn) limited (Browning et al., 2021; J. H. Martin et al., 1990), while most Arctic Ocean waters are believed to be largely N limited, with pockets of seasonal Fe limitation (Arrigo et al., 2017; Nielsdottir et al., 2009). The biogeochemical impact of direct micronutrient enrichment and indirect macronutrient enrichment is therefore complex and spatially variable, requiring more holistic studies in the future.

The meltwater pump provides a direct link between the benthic environment and the surface yet the role of benthic–pelagic coupling is poorly quantified but likely important. Diagenetic recycling of nutrients in coastal sediments with a high input of reactive glaciogenic sediment and organic matter is likely to be significant (e.g., Halbach et al., 2019; Hendry et al., 2019; Laufer-Meiser et al., 2021; Wehrmann et al., 2014), and the input of submarine groundwater in regions that are heavily glaciated is virtually unknown (Ó Dochartaigh et al., 2019). Sediment diagenesis and submarine groundwater discharge is potentially already imprinted on the biogeochemical data sets of previous studies yet is very hard to account for without direct observations of benthic processes in coastal sediments (Ng et al., 2020) coupled to numerical models (e.g., Hülse et al., 2018).

Linked to the benthic reprocessing of glacial flour is also the role of particle-bound phases in coastal nutrient cycling. Glacial meltwaters carry an abundance of reactive minerals that could supply macronutrients to biota after further biogeochemical processing (Hendry et al., 2019). Particle-bound nutrients are found at concentrations often over an order of magnitude higher than dissolved macronutrients in glacial meltwater rivers (J. R. Hawkings et al., 2015; Pryer et al., 2020), yet the fate of this material could be anywhere from burial/loss to remobilization, is temporally and spatially variable, and dependent on a range of external factors (e.g., organic matter reactivity and sedimentation rates). Glacial flour is highly bioavailable when used to relieve nutrient limitation in microcosm experiments (e.g., Shoenfelt et al., 2017) and particulate phases are dissolvable in seawater under lab conditions (e.g., J. R. Hawkings et al., 2017). However, the lability of these phases in the marine water column is not thought to be significant based on dissolved macronutrient data (Hopwood et al., 2020) and particles are unlikely to supply nitrogen, the primary limiting nutrient in Arctic marine systems (Bhatia et al., 2021; Randelhoff et al., 2020), in any significant quantity (Wadham et al., 2016). Much of the particulate flux is lost at low salinities as seawater-induced particle aggregation and flocculation accelerates sedimentation, and some evidence suggests adsorption of SRP onto reactive iron phases during seawater–freshwater mixing (Cape, Straneo, et al., 2019). Highly turbid waters also limit light penetration into the water column and provide a challenging environment for some eukaryotic organisms (Giesecke et al., 2019). The impact of particulate material (as nutrient and as a light modulator) is likely to be heavily dependent on the source geology (Halbach et al., 2019), which varies widely across glacially covered regions. The role of glacial flour on nutrient inventories therefore remains heavily debated.

Marine-terminating glaciers in the Arctic are rapidly retreating in response to atmospheric and/or oceanic warming (Cook et al., 2019; Wood et al., 2021). Glacier grounding lines are likely to retreat into shallower depths or onto a retrograde slope (and therefore deeper depths; e.g., in Greenland and Antarctica; Gudmundsson et al., 2012; King et al., 2020), the glacier may transition to a land-based system, and/or the subglacial hydrological system may expand or contract in response to external effects (e.g., meltwater input, ice thickness, and mean annual temperature). A transition from marine to land terminating system will almost certainly impact proximal ecosystems by eliminating the meltwater pump and increasing the discharge of turbid meltwater into the euphotic zone. Perhaps the ultimate question to pursue is therefore how glacial nutrient fluxes (direct and indirect) are likely to change under these different confounding scenarios.

The work of Williams et al. (2021) in the CAA highlights some of the nuances associated with understanding biogeochemical cycling at the ice–ocean interface. Studies of coastal polar processes need to acknowledge the complex interplay of physiochemical processes at this emerging critical zone (Oliver et al., 2018), with both known unknowns (e.g., benthic fluxes) and unknown unknowns. Collecting data in these environments is challenging due to the remoteness and associated logistical complexity, as well as the unpredictability of rapidly evolving landscapes. However, additional temporal and spatial coverage is needed to understand the evolution of these interfaces. Only with a better physical and biogeochemical understanding of these systems will we be able to accurately predict future changes to glaciomarine macronutrient cycling. Knowledge of the coupling between glacier nutrient supply and marine productivity in polar regions is essential because of the tight linkages with ocean–atmosphere CO₂ exchange and the health of marine ecosystems, both of which are highly relevant to society. The challenge is open to integrate observational data over ice to ocean gradients with more realistic numerical models that can be used to predict future response of these vulnerable systems in a warmer world.

Acknowledgments

I thank the Editor-in-Chief Marguerite Xenopoulos and Associate Editor Aaron Stubbins for their invitation to write this commentary. I am indebted to Kate Hendry, Martyn Tranter, and an anonymous reviewer for providing comments on this manuscript, which helped improve its quality. The title of this commentary was inspired by the paper “Trickle or treat: The dynamics of nutrient export from polar glaciers” by Ashley Dubnick and others.

Data Availability Statement

No new data was produced in this commentary.

References

- Aciego, S. M., Stevenson, E. I., & Arendt, C. A. (2015). Climate versus geological controls on glacial meltwater micronutrient production in southern Greenland. *Earth and Planetary Science Letters*, 424, 51–58. <https://doi.org/10.1016/j.epsl.2015.05.017>
- Anesio, A. M., Lutz, S., Christmas, N. A. M., & Benning, L. G. (2017). The microbiome of glaciers and ice sheets. *Npj Biofilms and Microbiomes*, 3(1), 10. <https://doi.org/10.1038/s41522-017-0019-0>
- Apolloni, S. (1973). Glaciers and nutrients in Arctic Seas. *Science*, 180(4085), 491–493. <https://doi.org/10.1126/science.180.4085.491>

- Arrigo, K. R., van Dijken, G. L., Castelain, R. M., Luo, H., Rennermalm, Å. K., Tedesco, M., et al. (2017). Melting glaciers stimulate large summer phytoplankton blooms in southwest Greenland waters. *Geophysical Research Letters*, *44*, 6278–6285. <https://doi.org/10.1002/2017GL073583>
- Bhatia, M. P., Kujawinski, E. B., Das, S. B., Breier, C. F., Henderson, P. B., & Charette, M. A. (2013). Greenland meltwater as a significant and potentially bioavailable source of iron to the ocean. *Nature Geoscience*, *6*(4), 274–278. <https://doi.org/10.1038/ngeo1746>
- Bhatia, M. P., Waterman, S., Burgess, D. O., Williams, P. L., Bundy, R. M., Mellett, T., et al. (2021). Glaciers and nutrients in the Canadian Arctic Archipelago marine system. *Global Biogeochemical Cycles*, *35*, e2021GB006976. <https://doi.org/10.1029/2021GB006976>
- Browning, T. J., Achterberg, E. P., Engel, A., & Mawji, E. (2021). Manganese co-limitation of phytoplankton growth and major nutrient drawdown in the Southern Ocean. *Nature Communications*, *12*(1), 884. <https://doi.org/10.1038/s41467-021-21122-6>
- Bruland, K. W., Rue, E. L., & Smith, G. J. (2001). Iron and macronutrients in California coastal upwelling regimes: Implications for diatom blooms. *Limnology & Oceanography*, *46*(7), 1661–1674. <https://doi.org/10.4319/lo.2001.46.7.1661>
- Burton, J. D., & Liss, P. S. (1973). Processes of supply and removal of dissolved silicon in the oceans. *Geochimica et Cosmochimica Acta*, *37*(7), 1761–1773. [https://doi.org/10.1016/0016-7037\(73\)90161-0](https://doi.org/10.1016/0016-7037(73)90161-0)
- Cape, M. R., Straneo, F., Beaird, N., Bundy, R. M., & Charette, M. A. (2019). Nutrient release to oceans from buoyancy-driven upwelling at Greenland tidewater glaciers. *Nature Geoscience*, *12*(1), 34–39. <https://doi.org/10.1038/s41561-018-0268-4>
- Cape, M. R., Vernet, M., Pettit, E. C., Wellner, J., Truffer, M., Akie, G., et al. (2019). Circumpolar deep water impacts glacial meltwater export and coastal biogeochemical cycling along the West Antarctic Peninsula. *Frontiers in Marine Science*, *6*, 144. <https://doi.org/10.3389/fmars.2019.00144>
- Chassot, E., Bonhommeau, S., Dulvy, N. K., Mélin, F., Watson, R., Gascuel, D., & Le Pape, O. (2010). Global marine primary production constrains fisheries catches. *Ecology Letters*, *13*(4), 495–505. <https://doi.org/10.1111/j.1461-0248.2010.01443.x>
- Chu, V. W. (2014). Greenland Ice Sheet hydrology: A review. *Progress in Physical Geography*, *38*(1), 19–54. <https://doi.org/10.1177/0309133313507075>
- Cook, A. J., Copland, L., Noël, B. P. Y., Stokes, C. R., Bentley, M. J., Sharp, M. J., et al. (2019). Atmospheric forcing of rapid marine-terminating glacier retreat in the Canadian Arctic Archipelago. *Science Advances*, *5*(3), eaau8507. <https://doi.org/10.1126/sciadv.aau8507>
- De La Rocha, C. L., & Passow, U. (2014). 8.4—The biological pump. In H. D. Holland & K. K. Turekian (Eds.), *Treatise on geochemistry* (2nd ed., pp. 93–122). Elsevier. <https://doi.org/10.1016/B978-0-08-095975-7.00604-5>
- Dubnick, A., Sharp, M., Danielson, B., Saidi-Mehrabad, A., & Barker, J. (2020). Basal thermal regime affects the biogeochemistry of subglacial systems. *Biogeosciences*, *17*(4), 963–977. <https://doi.org/10.5194/bg-17-963-2020>
- Fellman, J. B., Spencer, R. G. M., Hernes, P. J., Edwards, R. T., D'Amore, D. V., & Hood, E. (2010). The impact of glacier runoff on the biodegradability and biochemical composition of terrigenous dissolved organic matter in near-shore marine ecosystems. *Marine Chemistry*, *121*(1–4), 112–122. <https://doi.org/10.1016/J.Marchem.2010.03.009>
- Giesecke, R., Höfer, J., Vallejos, T., & González, H. E. (2019). Death in southern Patagonian fjords: Copepod community structure and mortality in land- and marine-terminating glacier-fjord systems. *Subantarctic and Antarctic Marine Ecosystems: Outlining Patterns and Processes in a Changing Ocean*, *174*, 162–172. <https://doi.org/10.1016/j.pocan.2018.10.011>
- Gove, J. M., McManus, M. A., Neuheimer, A. B., Polovina, J. J., Drazen, J. C., Smith, C. R., et al. (2016). Near-island biological hotspots in barren ocean basins. *Nature Communications*, *7*(1), 10581. <https://doi.org/10.1038/ncomms10581>
- Greisman, P. (1979). On upwelling driven by the melt of ice shelves and tidewater glaciers. *Deep Sea Research Part A. Oceanographic Research Papers*, *26*(9), 1051–1065. [https://doi.org/10.1016/0198-0149\(79\)90047-5](https://doi.org/10.1016/0198-0149(79)90047-5)
- Gudmundsson, G. H., Krug, J., Durand, G., Favier, L., & Gagliardini, O. (2012). The stability of grounding lines on retrograde slopes. *The Cryosphere*, *6*(6), 1497–1505. <https://doi.org/10.5194/tc-6-1497-2012>
- Halbach, L., Vihtakari, M., Duarte, P., Everett, A., Granskog, M. A., Hop, H., et al. (2019). Tidewater glaciers and bedrock characteristics control the phytoplankton growth environment in a Fjord in the Arctic. *Frontiers in Marine Science*, *6*, 254. <https://doi.org/10.3389/fmars.2019.00254>
- Hartley, C., & Dunbar, M. (1938). On the hydrographic mechanism of the so-called brown zones associated with tidal glaciers. *Journal of Marine Research*, *1*(4), 305–311.
- Hawkings, J., Wadham, J., Tranter, M., Telling, J., Bagshaw, E., Beaton, A., et al. (2016). The Greenland Ice Sheet as a hotspot of phosphorus weathering and export in the Arctic. *Global Biogeochemical Cycles*, *30*, 191–210. <https://doi.org/10.1002/2015GB005237>
- Hawkings, J. R., Skidmore, M. L., Wadham, J. L., Priscu, J. C., Morton, P. L., Hatton, J. E., et al. (2020). Enhanced trace element mobilization by Earth's ice sheets. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(50), 31648–31659. <https://doi.org/10.1073/pnas.2014378117>
- Hawkings, J. R., Wadham, J. L., Benning, L. G., Hendry, K. R., Tranter, M., Tedstone, A., et al. (2017). Ice sheets as a missing source of silica to the polar oceans. *Nature Communications*, *8*, 14198. <https://doi.org/10.1038/ncomms14198>
- Hawkings, J. R., Wadham, J. L., Tranter, M., Lawson, E., Sole, A., Cowton, T., et al. (2015). The effect of warming climate on nutrient and solute export from the Greenland Ice Sheet. *Geochemical Perspectives Letters*, *1*, 94–104. <https://doi.org/10.7185/geochemlet.1510>
- Hendry, K. R., Briggs, N., Henson, S., Opher, J., Brearley, J. A., Meredith, M. P., et al. (2021). Tracing glacial meltwater from the Greenland Ice Sheet to the ocean using gliders. *Journal of Geophysical Research: Oceans*, *126*, e2021JC017274. <https://doi.org/10.1029/2021JC017274>
- Hendry, K. R., Huvenne, V. A. I., Robinson, L. F., Annett, A., Badger, M., Jacobel, A. W., et al. (2019). The biogeochemical impact of glacial meltwater from southwest Greenland. *Progress in Oceanography*, *176*, 102126. <https://doi.org/10.1016/j.pocan.2019.102126>
- Herraiz-Borreguero, L., Lannuzel, D., van der Merwe, P., Treverrow, A., & Pedro, J. B. (2016). Large flux of iron from the Amery Ice Shelf marine ice to Prydz Bay, East Antarctica. *Journal of Geophysical Research: Oceans*, *121*, 6009–6020. <https://doi.org/10.1002/2016JC011687>
- Hood, E., & Scott, D. (2008). Riverine organic matter and nutrients in southeast Alaska affected by glacial coverage. *Nature Geoscience*, *1*(9), 583–587. <https://doi.org/10.1038/ngeo280>
- Hopwood, M. J., Carroll, D., Browning, T. J., Meire, L., Mortensen, J., Krisch, S., & Achterberg, E. P. (2018). Non-linear response of summertime marine productivity to increased meltwater discharge around Greenland. *Nature Communications*, *9*(1), 3256. <https://doi.org/10.1038/s41467-018-05488-8>
- Hopwood, M. J., Carroll, D., Dunse, T., Hodson, A., Holding, J. M., Iriarte, J. L., et al. (2020). Review article: How does glacier discharge affect marine biogeochemistry and primary production in the Arctic? *The Cryosphere*, *14*(4), 1347–1383. <https://doi.org/10.5194/tc-14-1347-2020>
- Hopwood, M. J., Connelly, D. P., Arendt, K. E., Juul-Pedersen, T., Stinchcombe, M. C., Meire, L., et al. (2016). Seasonal changes in Fe along a glaciated Greenlandic Fjord. *Frontiers of Earth Science*, *4*, 15. <https://doi.org/10.3389/feart.2016.00015>
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., et al. (2021). Accelerated global glacier mass loss in the early twenty-first century. *Nature*, *592*(7856), 726–731. <https://doi.org/10.1038/s41586-021-03436-z>

- Hülse, D., Arndt, S., Daines, S., Regnier, P., & Ridgwell, A. (2018). OMEN-SED 1.0: A novel, numerically efficient organic matter sediment diagenesis module for coupling to Earth system models. *Geoscientific Model Development*, 11(7), 2649–2689. <https://doi.org/10.5194/gmd-11-2649-2018>
- Irvine-Fynn, T. D. L., Hodson, A. J., Moorman, B. J., Vatne, G., & Hubbard, A. L. (2011). Polythermal glacier hydrology: A review. *Reviews of Geophysics*, 49, RG4002. <https://doi.org/10.1029/2010RG000350>
- Jickells, T. D. (1998). Nutrient biogeochemistry of the coastal zone. *Science*, 281(5374), 217–222. <https://doi.org/10.1126/science.281.5374.217>
- Jickells, T. D., Baker, A. R., & Chance, R. (2016). Atmospheric transport of trace elements and nutrients to the oceans. *Philosophical Transactions. Series A, Mathematical, Physical, and Engineering Sciences*, 374(2081), 20150286. <https://doi.org/10.1098/rsta.2015.0286>
- Kanna, N., Sugiyama, S., Fukamachi, Y., Nomura, D., & Nishioka, J. (2020). Iron supply by subglacial discharge into a fjord near the front of a marine-terminating glacier in northwestern Greenland. *Global Biogeochemical Cycles*, 34, e2020GB006567. <https://doi.org/10.1029/2020GB006567>
- Kanna, N., Sugiyama, S., Ohashi, Y., Sakakibara, D., Fukamachi, Y., & Nomura, D. (2018). Upwelling of macronutrients and dissolved inorganic carbon by a subglacial freshwater driven plume in Bowdoin Fjord, northwestern Greenland. *Journal of Geophysical Research: Biogeosciences*, 123, 1666–1682. <https://doi.org/10.1029/2017JG004248>
- Kellerman, A. M., Hawkings, J. R., Wadham, J. L., Kohler, T. J., Stibal, M., Grater, E., et al. (2020). Glacier outflow dissolved organic matter as a window into seasonally changing carbon sources: Leverett glacier, Greenland. *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005161. <https://doi.org/10.1029/2019JG005161>
- King, M. D., Howat, I. M., Candela, S. G., Noh, M. J., Jeong, S., Noël, B. P. Y., et al. (2020). Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat. *Communications Earth & Environment*, 1(1), 1. <https://doi.org/10.1038/s43247-020-0001-2>
- Krisch, S., Hopwood, M. J., Schaffer, J., Al-Hashem, A., Höfer, J., Rutgers van der Loeff, M. M., et al. (2021). The 79°N glacier cavity modulates subglacial iron export to the NE Greenland Shelf. *Nature Communications*, 12(1), 3030. <https://doi.org/10.1038/s41467-021-23093-0>
- Laidre, K. L., Moon, T., Hauser, D. D. W., McGovern, R., Heide-Jørgensen, M. P., Dietz, R., & Hudson, B. (2016). Use of glacial fronts by narwhals (*Monodon monoceros*) in West Greenland. *Biology Letters*, 12(10), 20160457. <https://doi.org/10.1098/rsbl.2016.0457>
- Laufer-Meisner, K., Michaud, A. B., Maisch, M., Byrne, J. M., Kappler, A., Patterson, M. O., et al. (2021). Potentially bioavailable iron produced through benthic cycling in glaciated Arctic fjords of Svalbard. *Nature Communications*, 12(1), 1349. <https://doi.org/10.1038/s41467-021-21558-w>
- Long, S., & Jones, P. J. S. (2021). Greenland's offshore Greenland halibut fishery and role of the Marine Stewardship Council certification: A governance case study. *Marine Policy*, 127, 104095. <https://doi.org/10.1016/j.marpol.2020.104095>
- Lydersen, C., Assmy, P., Falk-Petersen, S., Kohler, J., Kovacs, K. M., Reigstad, M., et al. (2014). The importance of tidewater glaciers for marine mammals and seabirds in Svalbard, Norway. *Journal of Marine Systems*, 129, 452–471. <https://doi.org/10.1016/j.jmarsys.2013.09.006>
- Martin, J. B., Pain, A. J., Martin, E. E., Rahman, S., & Ackerman, P. (2020). Comparisons of nutrients exported from Greenlandic glacial and deglaciated watersheds. *Global Biogeochemical Cycles*, 34, e2020GB006661. <https://doi.org/10.1029/2020GB006661>
- Martin, J. H., Fitzwater, S. E., & Gordon, R. M. (1990). Iron deficiency limits phytoplankton growth in Antarctic waters. *Global Biogeochemical Cycles*, 4(1), 5–12. <https://doi.org/10.1029/GB004i001p00005>
- McCutcheon, J., Lutz, S., Williamson, C., Cook, J. M., Tedstone, A. J., Vanderstraeten, A., et al. (2021). Mineral phosphorus drives glacier algal blooms on the Greenland Ice Sheet. *Nature Communications*, 12(1), 570. <https://doi.org/10.1038/s41467-020-20627-w>
- Meire, L., Meire, P., Struyf, E., Krawczyk, D. W., Arendt, K. E., Yde, J. C., et al. (2016). High export of dissolved silica from the Greenland Ice Sheet. *Geophysical Research Letters*, 43, 9173–9182. <https://doi.org/10.1002/2016GL070191>
- Meire, L., Mortensen, J., Meire, P., Juul-Pedersen, T., Sejr, M. K., Rysgaard, S., et al. (2017). Marine-terminating glaciers sustain high productivity in Greenland fjords. *Global Change Biology*, 23, 5344–5357. <https://doi.org/10.1111/gcb.13801>
- Moon, T., Sutherland, D. A., Carroll, D., Felikson, D., Kehrl, L., & Straneo, F. (2018). Subsurface iceberg melt key to Greenland fjord freshwater budget. *Nature Geoscience*, 11(1), 49–54. <https://doi.org/10.1038/s41561-017-0018-z>
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., et al. (2013). Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*, 6(9), 701–710. <https://doi.org/10.1038/ngeo1765>
- Mouginot, J., Rignot, E., Björk, A. A., van den Broeke, M., Millan, R., Morlighem, M., et al. (2019). Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018. *Proceedings of the National Academy of Sciences of the United States of America*, 116, 9239–9244. <https://doi.org/10.1073/pnas.1904242116>
- Ng, H. C., Cassarino, L., Pickering, R. A., Woodward, E. M. S., Hammond, S. J., & Hendry, K. R. (2020). Sediment efflux of silicon on the Greenland margin and implications for the marine silicon cycle. *Earth and Planetary Science Letters*, 529, 115877. <https://doi.org/10.1016/j.epsl.2019.115877>
- Nielsdóttir, M. C., Moore, C. M., Sanders, R., Hinz, D. J., & Achterberg, E. P. (2009). Iron limitation of the postbloom phytoplankton communities in the Iceland Basin. *Global Biogeochemical Cycles*, 23, GB3001. <https://doi.org/10.1029/2008GB003410>
- Nienow, P. W., Sole, A. J., Slater, D. A., & Cowton, T. R. (2017). Recent advances in our understanding of the role of meltwater in the Greenland Ice Sheet System. *Current Climate Change Reports*, 3(4), 330–344. <https://doi.org/10.1007/s40641-017-0083-9>
- Nishizawa, B., Kanna, N., Abe, Y., Ohashi, Y., Sakakibara, D., Asaji, I., et al. (2019). Contrasting assemblages of seabirds in the subglacial meltwater plume and oceanic water of Bowdoin Fjord, northwestern Greenland. *ICES Journal of Marine Science*, 77(2), 711–720. <https://doi.org/10.1093/icesjms/fsz213>
- Ó Dochartaigh, B. É., MacDonald, A. M., Black, A. R., Everest, J., Wilson, P., Darling, W. G., et al. (2019). Groundwater–glacier meltwater interaction in proglacial aquifers. *Hydrology and Earth System Sciences*, 23(11), 4527–4539. <https://doi.org/10.5194/hess-23-4527-2019>
- Oliver, H., Luo, H., Castela, R. M., van Dijken, G. L., Mattingly, K. S., Rosen, J. J., et al. (2018). Exploring the potential impact of Greenland meltwater on stratification, photosynthetically active radiation, and primary production in the Labrador Sea. *Journal of Geophysical Research: Oceans*, 123, 2570–2591. <https://doi.org/10.1002/2018JC013802>
- Pain, A. J., Martin, J. B., Martin, E. E., Rahman, S., & Ackermann, P. (2020). Differences in the quantity and quality of organic matter exported from Greenlandic glacial and deglaciated watersheds. *Global Biogeochemical Cycles*, 34, e2020GB006614. <https://doi.org/10.1029/2020GB006614>
- Pryer, H. V., Hawkings, J. R., Wadham, J. L., Robinson, L. F., Hendry, K. R., Hatton, J. E., et al. (2020). The influence of glacial cover on riverine silicon and iron exports in Chilean Patagonia. *Global Biogeochemical Cycles*, 34, e2020GB006611. <https://doi.org/10.1029/2020GB006611>
- Randelhoff, A., Holding, J., Janout, M., Sejr, M. K., Babin, M., Tremblay, J.-É., & Alkire, M. B. (2020). Pan-Arctic Ocean primary production constrained by turbulent nitrate fluxes. *Frontiers in Marine Science*, 7, 150. <https://doi.org/10.3389/fmars.2020.00150>
- Santos, I. R., Chen, X., Lecher, A. L., Sawyer, A. H., Moosdorf, N., Rodellas, V., et al. (2021). Submarine groundwater discharge impacts on coastal nutrient biogeochemistry. *Nature Reviews Earth & Environment*, 2, 307–323. <https://doi.org/10.1038/s43017-021-00152-0>

- Shoenfelt, E. M., Sun, J., Winckler, G., Kaplan, M. R., Borunda, A. L., Farrell, K. R., et al. (2017). High particulate iron(II) content in glacially sourced dusts enhances productivity of a model diatom. *Science Advances*, 3(6), e1700314. <https://doi.org/10.1126/sciadv.1700314>
- Smith, K. L., Sherman, A. D., Shaw, T. J., & Sprintall, J. (2013). Icebergs as unique Lagrangian ecosystems in polar seas. *Annual Review of Marine Science*, 5(5), 269–287. <https://doi.org/10.1146/annurev-marine-121211-172317>
- Statham, P. J. (2012). Nutrients in estuaries—An overview and the potential impacts of climate change. *The Science of the Total Environment*, 434(0), 213–227. <https://doi.org/10.1016/j.scitotenv.2011.09.088>
- Tranter, M., & Wadham, J. L. (2014). 7.5—Geochemical weathering in glacial and proglacial environments. In H. D. Holland & K. K. Turekian (Eds.), *Treatise on geochemistry* (2nd ed., pp. 157–173). Elsevier. <https://doi.org/10.1016/b978-0-08-095975-7.00505-2>
- Tseng, Y.-F., Lin, J., Dai, M., & Kao, S.-J. (2014). Joint effect of freshwater plume and coastal upwelling on phytoplankton growth off the Changjiang River. *Biogeosciences*, 11(2), 409–423. <https://doi.org/10.5194/bg-11-409-2014>
- Urbanski, J. A., Stempniewicz, L., Węslawski, J. M., Dragańska-Deja, K., Wochna, A., Goc, M., & Iliszko, L. (2017). Subglacial discharges create fluctuating foraging hotspots for sea birds in tidewater glacier bays. *Scientific Reports*, 7, 43999. <https://doi.org/10.1038/srep43999>
- Vick-Majors, T. J., Michaud, A. B., Skidmore, M. L., Turetta, C., Barbante, C., Christner, B. C., et al. (2020). Biogeochemical connectivity between freshwater ecosystems beneath the West Antarctic Ice Sheet and the sub-ice marine environment. *Global Biogeochemical Cycles*, 34, e2019GB006446. <https://doi.org/10.1029/2019GB006446>
- Wadham, J. L., Hawkings, J., Telling, J., Chandler, D., Alcock, J., O'Donnell, E., et al. (2016). Sources, cycling and export of nitrogen on the Greenland Ice Sheet. *Biogeosciences*, 13(22), 6339–6352. <https://doi.org/10.5194/bg-13-6339-2016>
- Wadham, J. L., Hawkings, J. R., Tarasov, L., Gregoire, L. J., Spencer, R. G. M., Gutjahr, M., et al. (2019). Ice sheets matter for the global carbon cycle. *Nature Communications*, 10(1), 3567. <https://doi.org/10.1038/s41467-019-11394-4>
- Watanabe, K., Kasai, A., Fukuzaki, K., Ueno, M., & Yamashita, Y. (2017). Estuarine circulation-driven entrainment of oceanic nutrients fuels coastal phytoplankton in an open coastal system in Japan. *Estuarine, Coastal and Shelf Science*, 184, 126–137. <https://doi.org/10.1016/j.ecss.2016.10.031>
- Wehrmann, L. M., Formolo, M. J., Owens, J. D., Raiswell, R., Ferdelman, T. G., Riedinger, N., & Lyons, T. W. (2014). Iron and manganese speciation and cycling in glacially influenced high-latitude fjord sediments (West Spitsbergen, Svalbard): Evidence for a benthic recycling-transport mechanism. *Geochimica et Cosmochimica Acta*, 141, 628–655. <https://doi.org/10.1016/j.gca.2014.06.007>
- Williams, P. L., Burgess, D. O., Waterman, S., Roberts, M., Bertrand, E. M., & Bhatia, M. P. (2021). Nutrient and carbon export from a tidewater glacier to the coastal ocean in the Canadian High Arctic Archipelago. *Journal of Geophysical Research: Biogeosciences*, 126, e2021JG006289. <https://doi.org/10.1029/2021JG006289>
- Wood, M., Rignot, E., Fenty, I., An, L., Björk, A., van den Broeke, M., et al. (2021). Ocean forcing drives glacier retreat in Greenland. *Science Advances*, 7(1), eaba7282. <https://doi.org/10.1126/sciadv.aba7282>
- Wyatt, F. R., & Sharp, M. J. (2015). Linking surface hydrology to flow regimes and patterns of velocity variability on Devon Ice Cap, Nunavut. *Journal of Glaciology*, 61(226), 387–399. <https://doi.org/10.3189/2015JG14J109>